



## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<p>(51) International Patent Classification <sup>6</sup> : A61K 31/38, 31/40, 31/44, 31/495, 31/505, C07D 209/30, 209/40, 213/75, 213/81, 213/82, 239/32, 241/18, 241/20, 333/06, 401/10, 401/12</p>	A1	<p>(11) International Publication Number: <b>WO 99/00126</b></p> <p>(43) International Publication Date: 7 January 1999 (07.01.99)</p>
<p>(21) International Application Number: PCT/US98/13384</p> <p>(22) International Filing Date: 26 June 1998 (26.06.98)</p> <p>(30) Priority Data: 60/050,881 26 June 1997 (26.06.97) US</p> <p>(71) Applicant (for all designated States except US): ELI LILLY AND COMPANY [US/US]; Lilly Corporate Center, Indianapolis, IN 46285 (US).</p> <p>(72) Inventors; and</p> <p>(75) Inventors/Applicants (for US only): BEIGHT, Douglas, Wade [US/US]; Apartment G, 9368 Benchmark Drive, Indianapolis, IN 46240 (US). CRAFT, Trelia, Joyce [US/US]; 10404 East 46th Street, Indianapolis, IN 46236 (US). FRANCISKOVICH, Jeffry, Bernard [US/US]; 5036 Quail Ridge Lane, Indianapolis, IN 46254 (US). GOODSON, Theodore, Junior [US/US]; 4045 Devon Drive, Indianapolis, IN 46226 (US). HALL, Steven, Edward [US/US]; 102 Nuttal Place, Chapel Hill, NC 27514 (US). HERRON, David, Kent [US/US]; 5945 Andover Road, Indianapolis, IN 46220 (US). KLIMKOWSKI, Valentine, Joseph [US/US]; 4504 Camelot Lane, Carmel, IN 46033 (US). KYLE, Jeffrey, Alan</p>		<p>[US/US]; 10434 Collingswood Lane, Fishers, IN 46038 (US). MASTERS, John, Joseph [US/US]; 8338 Crystal Pointe Lane, Indianapolis, IN 46236 (US). MENDEL, David [US/US]; 11348 Woods Bay Lane, Indianapolis, IN 46236 (US). MILOT, Guy [CA/US]; 2 Farrington Street, Foxborough, MA 02035 (US). SAWYER, Jason, Scott [US/US]; 5718 North Winthrop Avenue, Indianapolis, IN 46220 (US). SHUMAN, Robert, Theodore [US/US]; 830 Ashton Park Drive, Greenwood, IN 46143 (US). SMITH, Gerald, Floyd [US/US]; 825 Queenswood Court, Indianapolis, IN 46217 (US). TEBBE, Anne, Louise [US/US]; 6202 North Sherman Drive, Indianapolis, IN 46220 (US). TINSLEY, Jennifer, Marie [US/US]; 4542 State Road 39 North, Martinsville, IN 46151 (US). WEIR, Leonard, Crayton [US/US]; 6520 Englehardt Drive, Raleigh, NC 27613 (US). WIKEL, James, Howard [US/US]; 4068 Sunshine Way, Greenwood, IN 46142 (US). WILEY, Michael, Robert [US/US]; 7725 Langwood Drive, Indianapolis, IN 46268 (US). YEH, Ying, Kwong [US/US]; 5127 Briarstone Trace, Carmel, IN 46033 (US).</p> <p>(74) Agents: JACKSON, Thomas, E. et al.; Eli Lilly and Company, Lilly Corporate Center, Indianapolis, IN 46285 (US).</p> <p>(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, GM, GW, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).</p> <p><b>Published</b> With international search report.</p>
<p>(54) Title: ANTITHROMBOTIC AGENTS</p> <p>(57) Abstract</p> <p>This application relates to a compound of formula (I) (or a prodrug thereof or a pharmaceutically acceptable salt of the compound or prodrug thereof) as defined herein, pharmaceutical compositions thereof, and its use as an inhibitor of factor Xa, as well as a process for its preparation and intermediates therefor.</p>		

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	R	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

- 1 -

## ANTITHROMBOTIC AGENTS

This application claims the benefit of U.S. Provisional  
5 Application No. 60/050,881, filed June 26, 1997.

This invention relates to antithrombotic heterocycles  
which demonstrate activity as inhibitors of factor Xa and,  
accordingly, which are useful anticoagulants in mammals. In  
particular it relates to heterocycles having high  
10 anticoagulant activity, and antithrombotic activity. Thus,  
this invention relates to new inhibitors of factor Xa,  
pharmaceutical compositions containing the compounds as  
active ingredients, and the use of the compounds as  
anticoagulants for prophylaxis and treatment of  
15 thromboembolic disorders such as venous thrombosis,  
pulmonary embolism, arterial thrombosis, in particular  
myocardial ischemia, myocardial infarction and cerebral  
thrombosis, general hypercoagulable states and local  
hypercoagulable states, such as following angioplasty and  
20 coronary bypass operations, and generalized tissue injury as  
it relates to the inflammatory process. In addition, the  
antithrombotic agents are useful as anticoagulants in in  
vitro applications.

The process of blood coagulation, thrombosis, is  
25 triggered by a complex proteolytic cascade leading to the  
formation of thrombin. Thrombin proteolytically removes  
activation peptides from the A $\alpha$ -chains and the B $\beta$ -chains of  
fibrinogen, which is soluble in blood plasma, initiating  
insoluble fibrin formation. The formation of thrombin from  
30 prothrombin is catalyzed by factor Xa.

- 2 -

Anticoagulation currently is achieved by the administration of heparins and coumarins. Parenteral pharmacological control of coagulation and thrombosis is based on inhibition of thrombin through the use of heparins.

5 Heparins act indirectly on thrombin by accelerating the inhibitory effect of endogenous antithrombin III (the main physiological inhibitor of thrombin). Because antithrombin III levels vary in plasma and because clot-bound thrombin seems resistant to this indirect mechanism, heparins can be

10 an ineffective treatment. Because coagulation assays are believed to be associated with efficacy and with safety, heparin levels must be monitored with coagulation assays (particularly the activated partial thromboplastin time (APTT) assay). Coumarins impede the generation of thrombin

15 by blocking the posttranslational gamma-carboxylation in the synthesis of prothrombin and other proteins of this type. Because of their mechanism of action, the effect of coumarins can only develop slowly, 6-24 hours after administration. Further, they are not selective

20 anticoagulants. Coumarins also require monitoring with coagulation assays (particularly the prothrombin time (PT) assay).

Recently, interest has grown in small synthetic molecules which demonstrate potent direct inhibition of

25 thrombin and factor Xa. See, Jeremy J. Edmunds and Stephen T. Rapundalo (Annette M. Doherty, Section Editor), Annual Reports in Medicinal Chemistry, (1996), 31, 51-60.

Although the heparins and coumarins are effective anticoagulants, no commercial drug has yet emerged from the


30 small synthetic molecules; and despite the continuing promise for this class of compounds, there still exists a need for anticoagulants which act selectively on factor Xa or thrombin, and which, independent of antithrombin III, exert inhibitory action shortly after administration,

- 3 -

preferably by an oral route, and do not interfere with lysis of blood clots, as required to maintain homeostasis.

The present invention is directed to the discovery that the compounds of the present invention, as defined below,  
 5 are potent inhibitors of factor Xa which may have high bioavailability following oral administration.

According to the invention there is provided a method of inhibiting factor Xa comprising using an effective amount of a factor Xa inhibiting compound of formula I  
 10



wherein

- 15 A<sup>3</sup>, A<sup>4</sup>, A<sup>5</sup> and A<sup>6</sup>, together with the two carbons to which they are attached, complete a substituted heteroaromatic ring in which
- (a) one of A<sup>3</sup>, A<sup>4</sup>, A<sup>5</sup> and A<sup>6</sup> is N, and each of the others is CR<sup>3</sup>, CR<sup>4</sup>, CR<sup>5</sup> or CR<sup>6</sup>, respectively;
  - (b) two adjacent residues of A<sup>3</sup>, A<sup>4</sup>, A<sup>5</sup> and A<sup>6</sup> together  
 20 form S, and each of the others is CR<sup>3</sup>, CR<sup>4</sup>, CR<sup>5</sup> or CR<sup>6</sup>, respectively;
  - (c) two non-adjacent residues of A<sup>3</sup>, A<sup>4</sup>, A<sup>5</sup> and A<sup>6</sup> are each N, and each of the others is CR<sup>3</sup>, CR<sup>4</sup>, CR<sup>5</sup> or CR<sup>6</sup>, respectively; or
  - 25 (d) A<sup>3</sup> and A<sup>4</sup> together form a fused benz ring, and A<sup>5</sup> and A<sup>6</sup> together form -NH-;


wherein

- each of R<sup>3</sup>, R<sup>4</sup>, R<sup>5</sup> and R<sup>6</sup> is hydrogen, or one or two of R<sup>3</sup>, R<sup>4</sup>, R<sup>5</sup> and R<sup>6</sup> is independently chloro, bromo or methyl  
 30 and the others are hydrogen;

L<sup>1</sup> is -NH-CO- or -CO-NH- such that -L<sup>1</sup>-Q<sup>1</sup> is -NH-CO-Q<sup>1</sup> or -CO-NH-Q<sup>1</sup>;

- 4 -


$Q^1$  is phenyl, 2-furanyl, 2-thienyl, 4-thiazolyl,  
 2-pyridyl, 2-naphthyl, 1,2-dihydrobenzofuran-5-yl,  
 1,2-dihydrobenzofuran-6-yl or 1,2-benzisoxazol-6-yl in which  
 the phenyl may bear a 2-fluoro substituent or may bear one,  
 5 two or three substituents at the 3-, 4- or 5-position(s)  
 independently selected from halo, cyano, carbamoyl,  
 aminomethyl, methyl, methoxy, difluoromethoxy,  
 hydroxymethyl, formyl, vinyl, amino, hydroxy and  
 3,4-methylenedioxy, the 2-furanyl or 2-thienyl may bear a  
 10 chloro or methyl substituent at the 5-position, the  
 4-thiazolyl may bear an amino substituent at the 2-position,  
 the 2-pyridyl may bear an amino substituent at the  
 6-position, and the 1,2-benzisoxazol-6-yl may bear a chloro  
 or methyl substituent at the 3-position; or  $-CO-Q^1$  is  
 15 cyclopentenylcarbonyl or cyclohexenylcarbonyl;  
 $R^2$  is  $-L^{2A}-Q^{2A}$ ,  $-L^{2B}-Q^{2B}$ ,  $-L^{2C}-Q^{2C}$  or  $-L^{2D}-Q^{2D}$  wherein  
 $L^{2A}$  is a direct bond; and  
 $Q^{2A}$  is



20 in which D is carbonyl or  $-CHR^k-$  in which  $R^k$  is hydrogen,  
 hydroxy, (1-6C)alkoxy or  $-CH_2-R^j$  in which  $R^j$  is carboxy,  
 [(1-4C)alkoxy]carbonyl or carbamoyl which may bear one or  
 two (1-2C)alkyl substituents on the nitrogen; and one of  $R^m$   
 25 and  $R^n$  is hydrogen and the other is amino, bromo,  
 (1-4C)alkyl or (1-4C)alkoxy, or  $R^m$  and  $R^n$  together form a  
 benz ring;

$L^{2B}$  is  $-NH-CO-$ ,  $-O-CO-$ ,  $-CH_2-O-$  or  $-O-CH_2-$  such that  
 $-L^{2B}-Q^{2B}$  is  $-NH-CO-Q^{2B}$ ,  $-O-CO-Q^{2B}$ ,  $-CH_2-O-Q^{2B}$  or  $-O-CH_2-Q^{2B}$ ;  
 30 and

- 5 -

Q<sup>2B</sup> is

in which R<sup>O</sup> is hydrogen, halo, (1-6C)alkyl, (1-4C)alkoxy,  
 5 benzyloxy or (1-4C)alkylthio; and R<sup>P</sup> is 1-hydroxyethyl,  
 1-hydroxy-1-methylethyl, 1-methoxy-1-methylethyl,  
 4-piperidinyl, 4-pyridinyl, dimethylaminosulfonyl or -J-R<sup>Q</sup>  
 in which J is a single bond, methylene, carbonyl, oxo,  
 -S(O)<sub>q</sub>- (wherein q is 0, 1 or 2), or -NR<sup>r</sup>- (wherein R<sup>r</sup> is  
 10 hydrogen or methyl); and R<sup>Q</sup> is (1-6C)alkyl, phenyl,  
 3-pyridyl or 4-pyridyl;

L<sup>2C</sup> is -NR<sup>V</sup>-CO-X-, -NR<sup>V</sup>-CS-Y-, -CH<sub>2</sub>-CO-NR<sup>W</sup>-CH<sub>2</sub>-,  
 -O-CO-, -O-CH<sub>2</sub>-, -S-CH<sub>2</sub>- or -CH<sub>2</sub>-NR<sup>X</sup>-CH<sub>2</sub>- such that -L<sup>2C</sup>-Q<sup>2C</sup>  
 is -NR<sup>V</sup>-CO-X-Q<sup>2C</sup>, -NR<sup>V</sup>-CS-Y-Q<sup>2C</sup>, -CH<sub>2</sub>-CO-NR<sup>W</sup>-CH<sub>2</sub>-Q<sup>2C</sup>,  
 15 -O-CO-Q<sup>2C</sup>, -O-CH<sub>2</sub>-Q<sup>2C</sup>, -S-CH<sub>2</sub>-Q<sup>2C</sup> or -CH<sub>2</sub>-NR<sup>X</sup>-CH<sub>2</sub>-Q<sup>2C</sup> in  
 which X is -(CH<sub>2</sub>)<sub>x</sub>- (wherein x is 0, 1 or 2), -NR<sup>W</sup>-,  
 -NR<sup>W</sup>-CH<sub>2</sub>-, -O-, -O-CH<sub>2</sub>- or -S-CH<sub>2</sub>-; Y is -NR<sup>W</sup>-CH<sub>2</sub>- or  
 -O-CH<sub>2</sub>-; each of R<sup>V</sup> and R<sup>W</sup> is independently hydrogen, benzyl  
 or (1-6C)alkyl which is not branched at the α-position; and  
 20 R<sup>X</sup> is hydrogen, benzyloxycarbonyl or [(1-4C)alkoxy]carbonyl;  
 and

Q<sup>2C</sup> is 1-(4-pyridyl)piperidin-4-yl, 1-(4-pyridyl)-  
 piperidin-3-yl or 1-(4-pyridyl)pyrrolidin-3-yl in which the  
 pyridyl may bear a substituent at its 2-position selected  
 25 from cyano, aminomethyl, carboxy, hydroxymethyl and  
 (1-2C)alkyl;

L<sup>2D</sup> is -NH-CO- such that -L<sup>2D</sup>-Q<sup>2D</sup> is -NH-CO-Q<sup>2D</sup>; and  
 Q<sup>2D</sup> is selected from 4-(4-pyridinyl)benzyloxy, 9-oxo-  
 9H-fluoren-3-yl, benzo[b]thiophen-2-yl (which may bear a  
 30 chloro, methyl or methoxy substituent), benzofuran-2-yl  
 (which may bear a chloro, methyl or methoxy substituent),  
 4-(4-morpholinyl)-4-oxobutyl, and 4-piperidinyl or

- 6 -

3,4-didehydropiperidin-4-yl (either one bearing a substituent at the 1-position selected from methylsulfonyl, phenylsulfonyl, (1-5C)alkyl, (4-7C)cycloalkyl, tetrahydropyran-4-yl, 4-thiacyclohexyl and  $-\text{CH}_2-\text{R}^2$  in which  $\text{R}^2$  is  
5 isopropyl, cyclopropyl, phenyl, furyl, thienyl, 2-thiazolyl, or pyridyl in which the phenyl may bear one or two substituents independently selected from halo, cyano, hydroxy, methoxy, acetoxy, benzyloxy, amino, acetylamino, nitro and 3,4-methylenedioxy, and the thienyl or furyl may  
10 bear a methyl or nitro substituent);

or a prodrug of the compound of formula I;

or a pharmaceutically acceptable salt of the compound of formula I or prodrug thereof.

A particular factor Xa inhibiting compound of formula I  
15 is one wherein

$\text{A}^3$ ,  $\text{A}^4$ ,  $\text{A}^5$  and  $\text{A}^6$ , together with the two carbons to which they are attached, complete a substituted heteroaromatic ring in which

(a) one of  $\text{A}^3$ ,  $\text{A}^4$ ,  $\text{A}^5$  and  $\text{A}^6$  is N, and each of the others  
20 is  $\text{CR}^3$ ,  $\text{CR}^4$ ,  $\text{CR}^5$  or  $\text{CR}^6$ , respectively;

(b) two adjacent residues of  $\text{A}^3$ ,  $\text{A}^4$ ,  $\text{A}^5$  and  $\text{A}^6$  together form S, and each of the others is  $\text{CR}^3$ ,  $\text{CR}^4$ ,  $\text{CR}^5$  or  $\text{CR}^6$ , respectively;

(c) two non-adjacent residues of  $\text{A}^3$ ,  $\text{A}^4$ ,  $\text{A}^5$  and  $\text{A}^6$  are each  
25 N, and each of the others is  $\text{CR}^3$ ,  $\text{CR}^4$ ,  $\text{CR}^5$  or  $\text{CR}^6$ , respectively; or

(d)  $\text{A}^3$  and  $\text{A}^4$  together form a fused benz ring, and  $\text{A}^5$  and  $\text{A}^6$  together form  $-\text{NH}-$ ;

wherein

30 each of  $\text{R}^3$ ,  $\text{R}^4$ ,  $\text{R}^5$  and  $\text{R}^6$  is hydrogen, or one or two of  $\text{R}^3$ ,  $\text{R}^4$ ,  $\text{R}^5$  and  $\text{R}^6$  is independently chloro, bromo or methyl and the others are hydrogen;


$\text{L}^1$  is  $-\text{NH}-\text{CO}-$  or  $-\text{CO}-\text{NH}-$  such that  $-\text{L}^1-\text{Q}^1$  is  $-\text{NH}-\text{CO}-\text{Q}^1$  or  $-\text{CO}-\text{NH}-\text{Q}^1$ ;



- 7 -

Q<sup>1</sup> is phenyl, 2-thienyl, 4-thiazolyl, 2-pyridyl, 2-naphthyl or 1,2-benzisoxazol-6-yl in which the phenyl may bear one, two or three substituents at the 3-, 4- or 5-position(s) independently selected from halo, cyano, carbamoyl, aminomethyl, methyl, methoxy, hydroxymethyl, formyl, vinyl, amino, hydroxy and 3,4-methylenedioxy, the 2-thienyl may bear a chloro or methyl substituent at the 5-position, the 4-thiazolyl may bear an amino substituent at the 2-position, the 2-pyridyl may bear an amino substituent at the 6-position, and the 1,2-benzisoxazol-6-yl may bear a chloro or methyl substituent at the 3-position;

R<sup>2</sup> is -L<sup>2A</sup>-Q<sup>2A</sup>, -L<sup>2B</sup>-Q<sup>2B</sup>, -L<sup>2C</sup>-Q<sup>2C</sup> or -L<sup>2D</sup>-Q<sup>2D</sup> wherein L<sup>2A</sup> is a direct bond; and Q<sup>2A</sup> is




15

in which D is carbonyl or -CHR<sup>k</sup>- in which R<sup>k</sup> is hydrogen, hydroxy, (1-6C)alkoxy or -CH<sub>2</sub>-R<sup>j</sup> in which R<sup>j</sup> is carboxy, [(1-4C)alkoxy]carbonyl or carbamoyl which may bear one or two (1-2C)alkyl substituents on the nitrogen; and one of R<sup>m</sup> and R<sup>n</sup> is hydrogen and the other is amino, bromo, (1-4C)alkyl or (1-4C)alkoxy, or R<sup>m</sup> and R<sup>n</sup> together form a benz ring;

L<sup>2B</sup> is -NH-CO-, -O-CO-, -CH<sub>2</sub>-O- or -O-CH<sub>2</sub>- such that -L<sup>2B</sup>-Q<sup>2B</sup> is -NH-CO-Q<sup>2B</sup>, -O-CO-Q<sup>2B</sup>, -CH<sub>2</sub>-O-Q<sup>2B</sup> or -O-CH<sub>2</sub>-Q<sup>2B</sup>; and

Q<sup>2B</sup> is



- 8 -

- in which  $R^O$  is hydrogen, halo, (1-6C)alkyl, (1-4C)alkoxy, benzyloxy or (1-4C)alkylthio; and  $R^P$  is 1-hydroxyethyl, 1-hydroxy-1-methylethyl, 1-methoxy-1-methylethyl,
- 5 4-piperidinyl, 4-pyridinyl, dimethylaminosulfonyl or  $-J-R^Q$  in which J is a single bond, methylene, carbonyl, oxo,  $-S(O)_q-$  (wherein q is 0, 1 or 2), or  $-NR^I-$  (wherein  $R^I$  is hydrogen or methyl); and  $R^Q$  is (1-6C)alkyl, phenyl, 3-pyridyl or 4-pyridyl;
- 10  $L^{2C}$  is  $-NR^V-CO-X-$ ,  $-NR^V-CS-Y-$ ,  $-CH_2-CO-NR^W-CH_2-$ ,  $-O-CO-$ ,  $-O-CH_2-$ ,  $-S-CH_2-$  or  $-CH_2-NR^X-CH_2-$  such that  $-L^{2C}-Q^{2C}$  is  $-NR^V-CO-X-Q^{2C}$ ,  $-NR^V-CS-Y-Q^{2C}$ ,  $-CH_2-CO-NR^W-CH_2-Q^{2C}$ ,  $-O-CO-Q^{2C}$ ,  $-O-CH_2-Q^{2C}$ ,  $-S-CH_2-Q^{2C}$  or  $-CH_2-NR^X-CH_2-Q^{2C}$  in which X is  $-(CH_2)_x-$  (wherein x is 0, 1 or 2),  $-NR^W-CH_2-$ ,
- 15  $-O-CH_2-$  or  $-S-CH_2-$ ; Y is  $-NR^W-CH_2-$  or  $-O-CH_2-$ ; each of  $R^V$  and  $R^W$  is independently hydrogen, benzyl or (1-6C)alkyl which is not branched at the  $\alpha$ -position; and  $R^X$  is hydrogen, benzyloxycarbonyl or [(1-4C)alkoxy]carbonyl; and
- $Q^{2C}$  is 1-(4-pyridyl)piperidin-4-yl in which the pyridyl
- 20 may bear a substituent at its 2-position selected from cyano, aminomethyl, carboxy, hydroxymethyl and (1-2C)alkyl;
- $L^{2D}$  is  $-NH-CO-$  such that  $-L^{2D}-Q^{2D}$  is  $-NH-CO-Q^{2D}$ ; and
- $Q^{2D}$  is selected from 4-(4-pyridinyl)benzyloxy, 9-oxo-9H-fluoren-3-yl, benzo[b]thiophen-2-yl (which may bear a
- 25 chloro, methyl or methoxy substituent), benzofuran-2-yl (which may bear a chloro, methyl or methoxy substituent), 4-(4-morpholinyl)-4-oxobutyl, and 4-piperidinyl bearing a substituent at the 1-position selected from methylsulfonyl, phenylsulfonyl and  $-CH_2-R^Z$  in which  $R^Z$  is isopropyl,
- 30 cyclopropyl, phenyl, furyl, thienyl, 2-thiazolyl, or pyridyl in which the phenyl may bear one or two substituents independently selected from halo, cyano, hydroxy, methoxy, acetoxy, benzyloxy, amino, acetylamino, nitro and

- 9 -

3,4-methylenedioxy, and the thienyl or furyl may bear a methyl or nitro substituent;

or a prodrug of the compound of formula I;

or a pharmaceutically acceptable salt of the compound  
5 of formula I or prodrug thereof.

In addition, there is provided the use of a factor Xa inhibiting compound of formula I (or prodrug or salt) as described herein as an active ingredient in the manufacture of a medicament for use in producing an anticoagulant or  
10 antithrombotic effect.

The present invention also provides a method of inhibiting coagulation in a mammal comprising administering to a mammal in need of treatment, a coagulation inhibiting dose of a factor Xa inhibiting compound of formula I having  
15 any of the definitions herein.

The present invention further provides a method of inhibiting factor Xa comprising administering to a mammal in need of treatment, a factor Xa inhibiting dose of a factor Xa inhibiting compound of formula I having any of the  
20 definitions herein.

Further, the present invention provides a method of treating a thromboembolic disorder comprising administering to a mammal in need of treatment, an effective dose of a factor Xa inhibiting compound of formula I having any of the  
25 definitions herein.

In addition, there is provided the use of a factor Xa inhibiting compound of formula I having any of the definitions herein for the manufacture of a medicament for treatment of a thromboembolic disorder.

30 As an additional feature of the invention there is provided a pharmaceutical formulation comprising in association with a pharmaceutically acceptable carrier, diluent or excipient, a prodrug of a factor Xa inhibiting

- 10 -

compound of formula I (or of a pharmaceutically acceptable salt thereof) as provided in any of the descriptions herein.

In general, the factor Xa inhibiting compounds of formula I are believed to be novel and, thus, to constitute an additional aspect of the invention. Thus, according to the invention there is provided a novel compound of formula I (or a pharmaceutically acceptable salt thereof) according to any of the definitions herein of a compound of formula I, provided that the compound is not one which is not novel.

A pharmaceutically acceptable salt of an antithrombotic agent of the instant invention includes one which is an acid-addition salt made from a basic compound of formula I and an acid which provides a pharmaceutically acceptable anion, as well as a salt which is made from an acidic compound of formula I and a base which provides a pharmaceutically acceptable cation. Thus, a salt of a novel compound of formula I as provided herein made with an acid or base which affords a pharmaceutically acceptable counterion provides a particular aspect of the invention. Examples of such acids and bases are provided hereinbelow.

As an additional aspect of the invention there is provided a pharmaceutical formulation comprising in association with a pharmaceutically acceptable carrier, diluent or excipient, a novel compound of formula I (or a pharmaceutically acceptable salt thereof) as provided in any of the descriptions herein.

In this specification, the following definitions are used, unless otherwise described: Halo is fluoro, chloro, bromo or iodo. Alkyl, alkoxy, etc. denote both straight and branched groups; but reference to an individual radical such as "propyl" embraces only the straight chain ("normal") radical, a branched chain isomer such as "isopropyl" being specifically denoted. When two adjacent residues form a

- 11 -

(fused) benz ring, they form a cis,cis-buta-1,3-dien-1,4-diyl divalent radical.

It will be appreciated that certain compounds of formula I (or salts or prodrugs, etc.) may exist in, and be isolated in, isomeric forms, including tautomeric forms, cis- or trans-isomers, as well as optically active, racemic, or diastereomeric forms. It is to be understood that the present invention encompasses a compound of formula I in any of the tautomeric forms or as an a mixture thereof; or as a mixture of diastereomers, as well as in the form of an individual diastereomer, and that the present invention encompasses a compound of formula I as a mixture of enantiomers, as well as in the form of an individual enantiomer, any of which mixtures or form possesses inhibitory properties against factor Xa, it being well known in the art how to prepare or isolate particular forms and how to determine inhibitory properties against factor Xa by standard tests including those described below.

In addition, a compound of formula I (or salt or prodrug, etc.) may exhibit polymorphism or may form a solvate with water or an organic solvent. The present invention also encompasses any such polymorphic form, any solvate or any mixture thereof.

Particular values are listed below for radicals, substituents, and ranges, for illustration only, and they do not exclude other defined values or other values within defined ranges for the radicals and substituents.

For an alkyl group or the alkyl portion of an alkyl containing group such as, for example alkoxy, a particular value for (1-2C)alkyl is methyl or ethyl, and more particularly is methyl; for (1-4C)alkyl is methyl, ethyl, propyl, isopropyl, butyl, isobutyl, or t-butyl, and more particularly is methyl, isopropyl, butyl or t-butyl; for (1-6C)alkyl is methyl, ethyl, propyl, butyl, pentyl or

- 12 -

hexyl, and more particularly is methyl, butyl or hexyl. A particular value for halo is bromo or chloro, and more particularly is chloro.


A particular value for  $R^2$  is one selected from  
 5  $-L^2A-Q^2A$ ,  $-NH-CO-Q^2B$ ,  $-NR^V-CO-X-Q^2C$ ,  $-NR^V-CS-Y-Q^2C$ , and  $-NH-CO-Q^2D$ .

One particular compound of formula I is a pyridine in which one of  $A^3$ ,  $A^4$ ,  $A^5$  and  $A^6$  is N, and each of the others is  $CR^3$ ,  $CR^4$ ,  $CR^5$  or  $CR^6$ , respectively.

10 Another particular compound of formula I is a thiophene in which the two adjacent residues  $A^5$  and  $A^6$  together form S, and  $A^3$  and  $A^4$  are  $CR^3$  and  $CR^4$ , respectively.

Another particular compound of formula I is an indole in which the two adjacent residues  $A^5$  and  $A^6$  together form  
 15  $-NH-$ , and  $A^3$  and  $A^4$  together form a fused benz ring.

A further particular compound of formula I is a pyridine of formula Ia



20 wherein  $A^4$  is N, and  $L^1$ ,  $Q^1$  and  $R^2$  have any of the values defined herein.

A particular value for  $Q^1$  is 4-methoxyphenyl.

A particular value for  $R^2$  is, for example, (4-t-butylbenzoyl)amino, (4-methoxybenzoyl)amino, [4-(4-pyridyl)-  
 25 benzoyl]amino or [1-(4-pyridyl)piperidin-4-yl]methoxycarbonylamino.

One particular compound of formula I as described herein is one in which  $L^1-Q^1$  is  $-NH-CO-Q^1$ .

Another particular compound of formula I as described  
 30 herein is one in which  $L^1-Q^1$  is  $-CO-NH-Q^1$ .

A prodrug of a compound of formula I may be one formed in a conventional manner with a functional group of the


- 13 -

compound, such as with an amino, hydroxy or carboxy group.

A compound of formula I may be prepared by processes which include processes known in the chemical art for the production of any known compounds of formula I or of  
 5 structurally analogous compounds or by a novel process described herein. A process for the preparation of a novel compound of formula I (or a pharmaceutically acceptable salt thereof), novel processes for the preparation of a compound of formula I and novel intermediates for the manufacture of  
 10 a compound of formula I as defined above provide further features of the invention and are illustrated by the following procedures in which the meanings of the generic radicals are as defined above, unless otherwise specified. It will be recognized that it may be preferred or necessary  
 15 to prepare a compound of formula I in which a functional group is protected using a conventional protecting group, then to remove the protecting group to provide the compound of formula I.

Thus, there is provided a process for preparing a novel  
 20 compound of formula I (or a pharmaceutically acceptable salt thereof) as provided in any of the above descriptions which is selected from any of those described in the examples, including the following.

(A) For a compound of formula I in which the linkage  
 25 of  $R^2$  to the ring terminates in  $-NH-CO-$ ,  $-NR^V-CO-$  or  $-NR^V-CS-$ , acylating an amine of formula II,



II


30 or a corresponding amine in which the nitrogen bears the group  $R^V$ , using a corresponding acid which terminates with the group  $HO-CO-$  or  $HO-CS-$ , or an activated derivative

- 14 -

thereof. Typical activated derivatives include the acid halides, activated esters, including 4-nitrophenyl esters and those derived from coupling reagents, as well as (when the product is a urea or thiourea) isocyanates and


5 isothiocyanates. It may be preferred to deprotonate the amine using a strong base in anhydrous conditions for the acylation reaction, for example as described in Example 2, Part C.

(B) For a compound of formula I in which  $-L^1-Q^1$  is  
10  $-NH-CO-Q^1$ , acylating an amine of formula III



using an acid of formula  $HO-CO-Q^1$ , or an activated  
15 derivative thereof. The conditions used may be similar to those of process (A), above.

(C) For a compound of formula I in which  $-L^1-Q^1$  is  
-CO-NH- $Q^1$  and  $R^2$  is of the form  $-NH-CO-Q^2$ , acylating an  
amine of formula  $H_2N-Q^1$  using a [1,3]oxazine of formula IV,  
20




wherein  $Q^2$  represents, for example,  $Q^{2B}$ ,  $Q^{2C}$  or  $Q^{2D}$ .

(D) For a compound of formula I in which  $R^2$  is  
25  $-L^{2A}-Q^{2A}$  and D is carbonyl, diacylating a compound of formula II using an anhydride of formula V.



- 15 -



V

Whereafter, for any of the above procedures, when a functional group is protected using a protecting group,  
5 removing the protecting group.

Whereafter, for any of the above procedures, when a pharmaceutically acceptable salt of a compound of formula I is required, it is obtained by reacting the basic form of a basic compound of formula I with an acid affording a  
10 physiologically acceptable counterion or the acidic form of an acidic compound of formula I with a base affording a physiologically acceptable counterion or by any other conventional procedure.

A novel intermediate or starting material compound such  
15 as, for example, a novel compound of formula II, III or IV, etc., provides a further aspect of the invention.

As mentioned above, a compound corresponding to a compound of formula I but in which a functional group is protected may serve as an intermediate for a compound of  
20 formula I. Accordingly, such a protected intermediate for a novel compound of formula I provides a further aspect of the invention. Protecting groups are well known in the art, for example as described in T.W. Greene and P.G.M. Wuts, "Protecting Groups in Organic Synthesis" (1991). Further,  
25 the protecting group may be a functionalized resin, for example as disclosed in H.V. Meyers, et al., Molecular Diversity, (1995), 1, 13-20.

As mentioned above, the invention includes a pharmaceutically acceptable salt of the factor Xa inhibiting  
30 compound defined by the above formula I. A basic compound

- 16 -

of this invention possesses one or more functional groups sufficiently basic to react with any of a number of inorganic and organic acids affording a physiologically acceptable counterion to form a pharmaceutically acceptable salt. Acids commonly employed to form pharmaceutically acceptable acid addition salts are inorganic acids such as hydrochloric acid, hydrobromic acid, hydroiodic acid, sulfuric acid, phosphoric acid, and the like, and organic acids such as p-toluenesulfonic acid, methanesulfonic acid, oxalic acid, p-bromobenzenesulfonic acid, carbonic acid, succinic acid, citric acid, benzoic acid, acetic acid, and the like. Examples of such pharmaceutically acceptable salts thus are the sulfate, pyrosulfate, bisulfate, sulfite, bisulfite, phosphate, monohydrogenphosphate, dihydrogenphosphate, metaphosphate, pyrophosphate, chloride, bromide, iodide, acetate, propionate, decanoate, caprylate, acrylate, formate, isobutyrate, caproate, heptanoate, propiolate, oxalate, malonate, succinate, suberate, sebacate, fumarate, maleate, butyne-1,4-dioate, hexyne-1,6-dioate, benzoate, chlorobenzoate, methylbenzoate, dinitrobenzoate, hydroxybenzoate, methoxybenzoate, phthalate, sulfonate, xylenesulfonate, phenylacetate, phenylpropionate, phenylbutyrate, citrate, lactate, gamma-hydroxybutyrate, glycollate, tartrate, methanesulfonate, propanesulfonate, naphthalene-1-sulfonate, naphthalene-2-sulfonate, mandelate, and the like. Preferred pharmaceutically acceptable acid addition salts include those formed with mineral acids such as hydrochloric acid, hydrobromic acid and sulfuric acid.

For a compound of formula I which bears an acidic moiety, such as a carboxy group, a pharmaceutically acceptable salt may be made with a base which affords a pharmaceutically acceptable cation, which includes alkali metal salts (especially sodium and potassium), alkaline

- 17 -

earth metal salts (especially calcium and magnesium), aluminum salts and ammonium salts, as well as salts made from physiologically acceptable organic bases such as triethylamine, morpholine, piperidine and triethanolamine.

5        If not commercially available, a necessary starting material for the preparation of a compound of formula I may be prepared by a procedure which is selected from standard techniques of organic chemistry, including aromatic and heteroaromatic substitution and transformation, from  
10 techniques which are analogous to the syntheses of known, structurally similar compounds, and techniques which are analogous to the above described procedures or procedures described in the Examples. It will be clear to one skilled in the art that a variety of sequences is available for the  
15 preparation of the starting materials. Starting materials which are novel provide another aspect of the invention.

      Selective methods of substitution, protection and deprotection are well known in the art for preparation of a compound such as one of formula II, III, IV or VI discussed  
20 above.

      Generally, a basic compound of the invention is isolated best in the form of an acid addition salt. A salt of a compound of formula I formed with an acid such as one of those mentioned above is useful as a pharmaceutically  
25 acceptable salt for administration of the antithrombotic agent and for preparation of a formulation of the agent. Other acid addition salts may be prepared and used in the isolation and purification of the compounds.

      As noted above, the optically active isomers and  
30 diastereomers of the compounds of formula I are also considered part of this invention. Such optically active isomers may be prepared from their respective optically active precursors by the procedures described above, or by resolving the racemic mixtures. This resolution can be

- 18 -

carried out by derivatization with a chiral reagent followed by chromatography or by repeated crystallization. Removal of the chiral auxiliary by standard methods affords substantially optically pure isomers of the compounds of the present invention or their precursors. Further details  
5 regarding resolutions can be obtained in Jacques, et al., Enantiomers, Racemates, and Resolutions, John Wiley & Sons, 1981.

The compounds of the invention are believed to  
10 selectively inhibit factor Xa over other proteinases and nonenzyme proteins involved in blood coagulation without appreciable interference with the body's natural clot lysing ability (the compounds have a low inhibitory effect on  
15 fibrinolysis). Further, such selectivity is believed to permit use with thrombolytic agents without substantial interference with thrombolysis and fibrinolysis.

The invention in one of its aspects provides a method of inhibiting factor Xa in mammals comprising administering to a mammal in need of treatment an effective (factor Xa  
20 inhibiting) dose of a compound of formula I.

In another of its aspects, the invention provides a method of treating a thromboembolic disorder comprising administering to a mammal in need of treatment an effective (thromboembolic disorder therapeutic and/or prophylactic  
25 amount) dose of a compound of formula I.

The invention in another of its aspects provides a method of inhibiting coagulation in a mammal comprising administering to a mammal in need of treatment an effective (coagulation inhibiting) dose of a compound of formula I.

30 The factor Xa inhibition, coagulation inhibition and thromboembolic disorder treatment contemplated by the present method includes both medical therapeutic and/or prophylactic treatment as appropriate.

- 19 -

In a further embodiment the invention relates to treatment, in a human or animal, of a condition where inhibition of factor Xa is required. The compounds of the invention are expected to be useful in mammals, including man, in treatment or prophylaxis of thrombosis and hypercoagulability in blood and tissues. Disorders in which the compounds have a potential utility are in treatment or prophylaxis of thrombosis and hypercoagulability in blood and tissues. Disorders in which the compounds have a potential utility, in treatment and/or prophylaxis, include venous thrombosis and pulmonary embolism, arterial thrombosis, such as in myocardial ischemia, myocardial infarction, unstable angina, thrombosis-based stroke and peripheral arterial thrombosis. Further, the compounds have expected utility in the treatment or prophylaxis of atherosclerotic disorders (diseases) such as coronary arterial disease, cerebral arterial disease and peripheral arterial disease. Further, the compounds are expected to be useful together with thrombolytics in myocardial infarction. Further, the compounds have expected utility in prophylaxis for reocclusion after thrombolysis, percutaneous transluminal angioplasty (PTCA) and coronary bypass operations. Further, the compounds have expected utility in prevention of rethrombosis after microsurgery. Further, the compounds are expected to be useful in anticoagulant treatment in connection with artificial organs and cardiac valves. Further, the compounds have expected utility in anticoagulant treatment in hemodialysis and disseminated intravascular coagulation. A further expected utility is in rinsing of catheters and mechanical devices used in patients *in vivo*, and as an anticoagulant for preservation of blood, plasma and other blood products *in vitro*. Still further, the compounds have expected utility in other diseases where blood coagulation could be a fundamental contributing

- 20 -

process or a source of secondary pathology, such as cancer, including metastasis, inflammatory diseases, including arthritis, and diabetes. The anti-coagulant compound is administered orally or parenterally, e.g. by intravenous  
5 infusion (iv), intramuscular injection (im) or subcutaneously (sc).

The specific dose of a compound administered according to this invention to obtain therapeutic and/or prophylactic effects will, of course, be determined by the particular  
10 circumstances surrounding the case, including, for example, the compound administered, the rate of administration, the route of administration, and the condition being treated.

A typical daily dose for each of the above utilities is between about 0.01 mg/kg and about 1000 mg/kg. The dose  
15 regimen may vary, e.g. for prophylactic use a single daily dose may be administered or multiple doses such as 3 or 5 times daily may be appropriate. In critical care situations a compound of the invention is administered by iv infusion at a rate between about 0.01 mg/kg/h and about 20 mg/kg/h  
20 and preferably between about 0.1 mg/kg/h and about 5 mg/kg/h.

The method of this invention also is practiced in conjunction with a clot lysing agent e.g. tissue plasminogen activator (t-PA), modified t-PA, streptokinase or urokinase.  
25 In cases when clot formation has occurred and an artery or vein is blocked, either partially or totally, a clot lysing agent is usually employed. A compound of the invention can be administered prior to or along with the lysing agent or subsequent to its use, and preferably further is  
30 administered along with aspirin to prevent the reoccurrence of clot formation.

The method of this invention is also practiced in conjunction with a platelet glycoprotein receptor (IIb/IIIa) antagonist, that inhibits platelet aggregation. A compound

- 21 -

of the invention can be administered prior to or along with the IIb/IIIa antagonist or subsequent to its use to prevent the occurrence or reoccurrence of clot formation.

The method of this invention is also practiced in conjunction with aspirin. A compound of the invention can be administered prior to or along with aspirin or subsequent to its use to prevent the occurrence or reoccurrence of clot formation. As stated above, preferably a compound of the present invention is administered in conjunction with a clot  
5  
10 lysing agent and aspirin.

This invention also provides a pharmaceutical composition for use in the above described therapeutic method. A pharmaceutical composition of the invention comprises an effective factor Xa inhibiting amount of a compound of formula I in association with a pharmaceutically  
15 acceptable carrier, excipient or diluent.

The active ingredient in such formulations comprises from 0.1 percent to 99.9 percent by weight of the formulation. By "pharmaceutically acceptable" it is meant the carrier, diluent or excipient must be compatible with  
20 the other ingredients of the formulation and not deleterious to the recipient thereof.

For oral administration the antithrombotic compound is formulated in gelatin capsules or tablets which may contain excipients such as binders, lubricants, disintegration  
25 agents and the like. For parenteral administration the antithrombotic is formulated in a pharmaceutically acceptable diluent e.g. physiological saline (0.9 percent), 5 percent dextrose, Ringer's solution and the like.

The compound of the present invention can be formulated in unit dosage formulations comprising a dose between about 0.1 mg and about 1000 mg. Preferably the compound is in the form of a pharmaceutically acceptable salt such as for  
30 example the sulfate salt, acetate salt or a phosphate salt.

- 22 -

An example of a unit dosage formulation comprises 5 mg of a compound of the present invention as a pharmaceutically acceptable salt in a 10 mL sterile glass ampoule. Another example of a unit dosage formulation comprises about 10 mg  
5 of a compound of the present invention as a pharmaceutically acceptable salt in 20 mL of isotonic saline contained in a sterile ampoule.

The compounds can be administered by a variety of routes including oral, rectal, transdermal, subcutaneous,  
10 intravenous, intramuscular, and intranasal. The compounds of the present invention are preferably formulated prior to administration.

The present pharmaceutical compositions are prepared by known procedures using well known and readily available  
15 ingredients. The compositions of this invention may be formulated so as to provide quick, sustained, or delayed release of the active ingredient after administration to the patient by employing procedures well known in the art. In making the compositions of the present invention, the active  
20 ingredient will usually be admixed with a carrier, or diluted by a carrier, or enclosed within a carrier which may be in the form of a capsule, sachet, paper or other container. When the carrier serves as a diluent, it may be a solid, semi-solid or liquid material which acts as a  
25 vehicle, excipient or medium for the active ingredient. Thus, the compositions can be in the form of tablets, pills, powders, lozenges, sachets, cachets, elixirs, suspensions, emulsions, solutions, syrups, aerosols, (as a solid or in a liquid medium), soft and hard gelatin capsules,  
30 suppositories, sterile injectable solutions, sterile packaged powders, and the like.

The following formulation examples are illustrative only and are not intended to limit the scope of the invention in any way. "Active ingredient," of course, means



- 23 -

a compound according to formula I or a pharmaceutically acceptable salt or solvate thereof.

- Formulation 1: Hard gelatin capsules are prepared  
5 using the following ingredients:

	Quantity (mg/capsule)
Active ingredient	250
Starch, dried	200
Magnesium stearate	<u>10</u>
Total	460 mg

- Formulation 2: A tablet is prepared using the ingredients below:

	Quantity (mg/tablet)
Active ingredient	250
Cellulose, microcrystalline	400
Silicon dioxide, fumed	10
Stearic acid	<u>5</u>
Total	665 mg

- 10 The components are blended and compressed to form tablets each weighing 665 mg.

- Formulation 3: An aerosol solution is prepared containing the following components:

	Weight
Active ingredient	0.25
Ethanol	29.75
Propellant 22 (Chlorodifluoromethane)	<u>70.00</u>
Total	100.00

- 24 -

The active compound is mixed with ethanol and the mixture added to a portion of the propellant 22, cooled to -30 °C and transferred to a filling device. The required amount is then fed to a stainless steel container and diluted with the remainder of the propellant. The valve units are then fitted to the container.

Formulation 4: Tablets, each containing 60 mg of active ingredient, are made as follows:

Active ingredient	60 mg
Starch	45 mg
Microcrystalline cellulose	35 mg
Polyvinylpyrrolidone (as 10% solution in water)	4 mg
Sodium carboxymethyl starch	4.5 mg
Magnesium stearate	0.5 mg
Talc	<u>1 mg</u>
Total	150 mg

The active ingredient, starch and cellulose are passed through a No. 45 mesh U.S. sieve and mixed thoroughly. The aqueous solution containing polyvinylpyrrolidone is mixed with the resultant powder, and the mixture then is passed through a No. 14 mesh U.S. sieve. The granules so produced are dried at 50 °C and passed through a No. 18 mesh U.S. Sieve. The sodium carboxymethyl starch, magnesium stearate and talc, previously passed through a No. 60 mesh U.S. sieve, are then added to the granules which, after mixing, are compressed on a tablet machine to yield tablets each weighing 150 mg.

- 25 -

Formulation 5: Capsules, each containing 80 mg of active ingredient, are made as follows:

Active ingredient	80 mg
Starch	59 mg
Microcrystalline cellulose	59 mg
Magnesium stearate	<u>2 mg</u>
Total	200 mg

- 5 The active ingredient, cellulose, starch, and magnesium stearate are blended, passed through a No. 45 mesh U.S. sieve, and filled into hard gelatin capsules in 200 mg quantities.

- 10 Formulation 6: Suppositories, each containing 225 mg of active ingredient, are made as follows:

Active ingredient	225 mg
Saturated fatty acid glycerides	<u>2,000 mg</u>
Total	2,225 mg

- 15 The active ingredient is passed through a No. 60 mesh U.S. sieve and suspended in the saturated fatty acid glycerides previously melted using the minimum heat necessary. The mixture is then poured into a suppository mold of nominal 2 g capacity and allowed to cool.

- 20 Formulation 7: Suspensions, each containing 50 mg of active ingredient per 5 mL dose, are made as follows:

Active ingredient	50 mg
Sodium carboxymethyl cellulose	50 mg
Syrup	1.25 mL
Benzoic acid solution	0.10 mL

- 26 -

Flavor	q.v.
Color	q.v.
Purified water to total	5 mL

The active ingredient is passed through a No. 45 mesh U.S. sieve and mixed with the sodium carboxymethyl cellulose and syrup to form a smooth paste. The benzoic acid solution, flavor and color are diluted with a portion of the water and added, with stirring. Sufficient water is then added to produce the required volume.

Formulation 8: An intravenous formulation may be prepared as follows:

Active ingredient	100 mg
Isotonic saline	1,000 mL

The solution of the above ingredients generally is administered intravenously to a subject at a rate of 1 mL per minute.

The ability of a compound of the present invention to be an effective and orally active factor Xa inhibitor may be evaluated in one or more of the following assays or in other standard assays known to those in the art.

The inhibition by a compound of the inhibition of a serine protease of the human blood coagulation system or of the fibrinolytic system, as well as of trypsin, is determined in vitro for the particular enzyme by measuring its inhibitor binding affinity in an assay in which the enzyme hydrolyzes a particular chromogenic substrate, for example as described in Smith, G.F.; Gifford-Moore, D.; Craft, T.J.; Chirgadze, N.; Ruterbories, K.J.; Lindstrom, T.D.; Satterwhite, J.H. *Efegatran: A New Cardiovascular Anticoagulant. New Anticoagulants for the Cardiovascular*

- 27 -

Patient; Pifarre, R., Ed.; Hanley & Belfus, Inc.:  
Philadelphia, 1997; pp. 265-300. The inhibitor binding  
affinity is measured as apparent association constant  $K_{ass}$   
which is the hypothetical equilibrium constant for the  
5 reaction between enzyme and the test inhibitor compound (I).



$$K_{ass} = \frac{[\text{Enzyme-I}]}{[(\text{Enzyme}) \times (\text{I})]}$$

Conveniently, enzyme inhibition kinetics are performed  
10 in 96-well polystyrene plates and reaction rates are  
determined from the rate of hydrolysis of appropriate  
p-nitroanilide substrates at 405 nm using a Thermomax plate  
reader from Molecular Devices (San Francisco, CA). The same  
protocol is followed for all enzymes studied: 50  $\mu\text{L}$  buffer  
15 (0.03 M Tris, 0.15 M NaCl pH 7) in each well, followed by  
25  $\mu\text{L}$  of inhibitor solution (in 100% methanol, or in 50% v:v  
aqueous methanol) and 25  $\mu\text{L}$  enzyme solution; within two  
minutes, 150  $\mu\text{L}$  aqueous solution of chromogenic substrate  
(0.25 mg/mL) is added to start the enzymatic reaction. The  
20 rates of chromogenic substrate hydrolysis reactions provide  
a linear relationship with the enzymes studied such that  
free enzyme can be quantitated in reaction mixtures. Data  
is analyzed directly as rates by the Softmax program to  
produce [free enzyme] calculations for tight-binding  $K_{ass}$   
25 determinations. For apparent  $K_{ass}$  determinations, 1.34 nM  
human factor Xa is used to hydrolyze 0.18 mM BzIle-Glu-Gly-  
Arg-pNA; 5.9 nM human thrombin or 1.4 nM bovine trypsin is  
used to hydrolyze 0.2 mM BzPhe-Val-Arg-pNA; 3.4 nM human  
plasmin is used with 0.5 mM HD-Val-Leu-Lys-pNA; 1.2 nM human  
30 nt-PA is used with 0.81 mM HD-Ile-Pro-Arg-pNA; and 0.37 nM  
urokinase is used with 0.30 mM pyro-gfsGlu-Gly-Arg-pNA.

- 28 -

Kass is calculated for a range of concentrations of test compounds and the mean value reported in units of liter per mole. In general, a factor Xa inhibiting compound of formula I of the instant invention exhibits a Kass of 0.1 to 5 0.5 x 10<sup>6</sup> L/mole or much greater.

The factor Xa inhibitor preferably should spare fibrinolysis induced by urokinase, tissue plasminogen activator (t-PA) and streptokinase. This would be important to the therapeutic use of such an agent as an adjunct to 10 streptokinase, tp-PA or urokinase thrombolytic therapy and to the use of such an agent as an endogenous fibrinolysis-sparing (with respect to t-PA and urokinase) antithrombotic agent. In addition to the lack of interference with the amidase activity of the fibrinolytic proteases, such 15 fibrinolytic system sparing can be studied by the use of human plasma clots and their lysis by the respective fibrinolytic plasminogen activators.

#### Materials

20 Dog plasma is obtained from conscious mixed-breed hounds (either sex Butler Farms, Clyde, New York, U.S.A.) by venipuncture into 3.8 percent citrate. Fibrinogen is prepared from fresh dog plasma and human fibrinogen is prepared from in-date ACD human blood at the fraction I-2 25 according to previous procedures and specification. Smith, Biochem. J., 185, 1-11 (1980; and Smith, et al., Biochemistry, 11, 2958-2967, (1972). Human fibrinogen (98 percent pure/plasmin free) is from American Diagnostica, Greenwich, Connecticut. Radiolabeling of fibrinogen I-2 30 preparations is performed as previously reported. Smith, et al., Biochemistry, 11, 2958-2967, (1972). Urokinase is purchased from Leo Pharmaceuticals, Denmark, as 2200 Ploug units/vial. Streptokinase is purchased from Hoechst-Roussel Pharmaceuticals, Somerville, New J rsey.

Thrombus Formation and Compound Administration Regimens

Electrolytic injury of the intima of the LCX is produced by applying 100- $\mu$ A direct current (DC) to the anode. The current is maintained for 60 min and then discontinued whether the vessel has occluded or not. Thrombus formation proceeds spontaneously until the LCX is totally occluded (determined as zero CBF and an increase in the S-T segment). Compound administration is started after the occluding thrombus is allowed to age for 1 hour. A 2-hour infusion of the compounds of the present invention at doses of 0.5 and 1 mg/kg/hour is begun simultaneously with an infusion of thrombolytic agent (e.g. tissue plasminogen activator, streptokinase, APSAC). Repetition is followed for 3 hour after administration of test compound. Reocclusion of coronary arteries after successful thrombolysis is defined as zero CBF which persisted for at least 30 minutes.

Hematology and template bleeding time determinations

Whole blood cell counts, hemoglobin, and hematocrit values are determined on a 40- $\mu$ L sample of citrated (3.8 percent) blood (1 part citrate:9 parts blood) with a hematology analyzer (Cell-Dyn 900, Sequoia-Turner, Mount View, CA, U.S.A.). Gingival template bleeding times are determined with a Simplate II bleeding time device (Organon Teknika Durham, N.C., U.S.A.). The device is used to make 2 horizontal incisions in the gingiva of either the upper or lower left jaw of the dog. Each incision is 3 mm wide x 2 mm deep. The incisions are made, and a stopwatch is used to determine how long bleeding occurs. A cotton swab is used to soak up the blood as it oozes from the incision. Template bleeding time is the time from incision to stoppage of bleeding. Bleeding times are taken just before administration of test compound (0 min), 60 min into infusion, at conclusion of administration of the test compound (120 min), and at the end of the experiment.





which means are different. Significance level for rejection of the null hypothesis of equal means is  $P < 0.05$ .

5 Animals  
Male dogs (Beagles; 18 months - 2 years; 12-13 kg, Marshall Farms, North Rose, New York 14516) are fasted overnight and fed Purina certified Prescription Diet (Purina Mills, St. Louis, Missouri) 240 minutes after dosing. Water is available *ad libitum*. The room temperature is maintained between 66-74 °F; 45-50 percent relative humidity; and lighted from 0600-1800 hours.

Pharmacokinetic model.  
15 Test compound is formulated immediately prior to dosing by dissolving in sterile 0.9 percent saline to a 5 mg/mL preparation. Dogs are given a single 2 mg/kg dose of test compound by oral gavage. Blood samples (4.5 mL) are taken from the cephalic vein at 0.25, 0.5, 0.75, 1, 2, 3, 4 and 6 hours after dosing. Samples are collected in citrated Vacutainer tubes and kept on ice prior to reduction to HPLC MS. Plasma concentration of test compound is recorded and used to calculate the pharmacokinetic parameters: elimination rate constant,  $K_e$ ; total clearance,  $CL$ ; volume of distribution,  $V_d$ ; time of maximum plasma test compound concentration,  $T_{max}$ ; maximum concentration of test compound of  $T_{max}$ ,  $C_{max}$ ; plasma half-life,  $t_{0.5}$ ; and area under the curve, A.U.C.; fraction of test compound absorbed,  $F$ .

30 Canine Model of Coronary Artery Thrombosis  
Surgical preparation and instrumentation of the dogs are as described in Jackson, et al., Circulation, 82, 930-940 (1990). Mixed-breed hounds (aged 6-7 months, either sex, Butler Farms, Clyde, New York, U.S.A.) are anesthetized with sodium pentobarbital (30 mg/kg intravenously, i.v.), intubated, and ventilated with room air. Tidal volume and



shunt circuit. Blood is circulated through the shunt for 15 min before the thread is carefully removed and weighed. The weight of a wet thread is subtracted from the total weight of the thread and thrombus (see J.R. Smith, *Br J Pharmacol*, 77:29, 1982).

#### Fec13 model of arterial injury

The carotid arteries are isolated via a midline ventral cervical incision. A thermocouple is placed under each artery and vessel temperature is recorded continuously on a strip chart recorder. A cuff of tubing (0.058 ID x 0.077 OD x 4 mm, Baxter Med. Grade Silicone), cut longitudinally, is placed around each carotid directly above the thermocouple. Fec13 hexahydrate is dissolved in water and the concentration (20 percent) is expressed in terms of the actual weight of Fec13 only. To injure the artery and induce thrombosis, 2.85  $\mu$ L is pipetted into the cuff to bathe the artery above the thermocouple probe. Arterial occlusion is indicated by a rapid drop in temperature. The time to occlusion is reported in minutes and represents the elapsed time between application of Fec13 and the rapid drop in vessel temperature (see K.D. Kurz, *Thromb. Res.*, 60:269, 1990).

#### Coagulation parameters

Plasma thrombin time (TT) and activated partial thromboplastin time (APTT) are measured with a fibrometer. Blood is sampled from a jugular catheter and collected in syringe containing sodium citrate (3.8 percent, 1 part to 9 parts blood). To measure TT, rat plasma (0.1 mL) is mixed with saline (0.1 mL) and bovine thrombin (0.1 mL, 30 U/mL in TRIS buffer; Parke Davis) at 37 °C. For APTT, plasma (0.1 mL) and APTT solution (0.1 mL, Organon Teknika) are incubated for 5 minutes (37 °C) and CaCl<sub>2</sub> (0.1 mL, 0.025 M) is added to start coagulation. Assays are done in duplicate and averaged.





- 37 -

Et<sub>3</sub>N = triethylamine  
 Et<sub>2</sub>O = diethyl ether  
 EtOH = ethanol  
 FAB = Fast Atom Bombardment (Mass Spectroscopy)  
 Hex = hexanes  
 HOAt = 1-hydroxy-7-azabenzotriazole  
 HOBt = 1-hydroxybenzotriazole  
 HPLC = High Performance Liquid Chromatography  
 HRMS = high resolution mass spectrum  
 i-PrOH = isopropanol  
 IR = Infrared Spectrum  
 Me = methyl  
 MeI = methyl iodide  
 MeOH = methanol  
 MS-FD = field desorption mass spectrum  
 NBS = N-bromosuccinimide  
 NMR = Nuclear Magnetic Resonance  
 Ph = phenyl  
 i-Pr = isopropyl  
 RP/HPLC = Reversed Phase High Performance Liquid  
 Chromatography

5

HOAt = 1-hydroxy-7-azabenzotriazole  
 HOBt = 1-hydroxybenzotriazole  
 HPLC = High Performance Liquid Chromatography  
 HRMS = high resolution mass spectrum  
 i-PrOH = isopropanol  
 IR = Infrared Spectrum  
 Me = methyl  
 MeI = methyl iodide  
 MeOH = methanol  
 MS-FD = field desorption mass spectrum  
 NBS = N-bromosuccinimide  
 NMR = Nuclear Magnetic Resonance  
 Ph = phenyl  
 i-Pr = isopropyl  
 RP/HPLC = Reversed Phase High Performance Liquid  
 Chromatography

10

IR = Infrared Spectrum  
 Me = methyl  
 MeI = methyl iodide  
 MeOH = methanol  
 MS-FD = field desorption mass spectrum  
 NBS = N-bromosuccinimide  
 NMR = Nuclear Magnetic Resonance  
 Ph = phenyl  
 i-Pr = isopropyl  
 RP/HPLC = Reversed Phase High Performance Liquid  
 Chromatography

15

RP/HPLC = Reversed Phase High Performance Liquid  
 Chromatography

20

satd = saturated  
 SiO<sub>2</sub> = silica gel  
 TBS = tert-butyldimethylsilyl  
 TFA = trifluoroacetic acid  
 THF = tetrahydrofuran  
 TIPS = trisopropylsilyl  
 TLC = thin layer chromatography  
 tosyl = p-toluenesulfonyl  
 triflic acid = trifluoromethanesulfonic acid

25

30

Unless otherwise stated, pH adjustments and work up are  
 with aqueous acid or base solutions. <sup>1</sup>H-NMR indicates a  
 satisfactory NMR spectrum was obtained for the compound  
 described. IR indicates a satisfactory infra red spectrum  
 was obtained for the compound described.

35



- 36 -

All data are analyzed by one-way analysis of variance (ANOVA) followed by Student-Neuman-Kuels post hoc t test to determine the level of significance. Repeated-measures ANOVA are used to determine significant differences between  
5 time points during the experiments. Values are determined to be statistically different at least at the level of  $p < 0.05$ . All values are mean  $\pm$  SEM. All studies are conducted in accordance with the guiding principles of the American Physiological Society. Further details regarding  
10 the procedures are described in Jackson, et al., J. Cardiovasc. Pharmacol., (1993), 21, 587-599.

The following Examples are provided to further describe the invention and are not to be construed as  
15 limitations thereof.

The abbreviations, symbols and terms used in the examples have the following meanings.


Ac = acetyl  
AIBN = azobisisobutyronitrile  
20 Anal. = elemental analysis  
aq = aqueous  
Bn or Bzl = benzyl  
Boc = t-butyloxycarbonyl  
Bu = butyl  
25 n-BuLi = butyllithium  
Calc = calculated  
conc = concentrated  
DCC = dicyclohexylcarbodiimide  
DMAP = 4-dimethylaminopyridine  
30 DMF = dimethylformamide  
DMSO = dimethylsulfoxide  
EDC = 1-(3-dimethylaminopropyl)-3-ethyl-carbodiimide hydrochloride  
eq = (molar) equivalent  
35 Et = ethyl  
EtOAc = ethyl acetate



- 39 -

## Example 1

Preparation of 3-(4-tert-butylbenzoyl)amino-N-(4-methoxyphenyl)-2-thiophenecarboxamide.



5

A. methyl 3-(4-tert-butylbenzoyl)amino-2-thiophenecarboxylate

A solution of methyl 3-amino-2-thiophenecarboxylate (400 mg, 2.54 mmol) and pyridine (0.226 mL, 2.80 mmol) in methylene chloride (12 mL) was treated with 4-tert-butylbenzoyl chloride (0.500 mL, 2.54 mmol). After consumption of the starting material, the mixture was concentrated in vacuo and the residue dissolved in ethyl acetate and water. The organic layer was washed once with water, once with saturated sodium chloride solution, dried (magnesium sulfate), and filtered. Concentration and purification of the residue by flash chromatography (silica gel, hexanes/ethyl acetate) yielded 689 mg (85%) of the title compound.

20

<sup>1</sup>H-NMR, IR

MS-FD m/e 317 (M<sup>+</sup>)

Analysis for C<sub>17</sub>H<sub>19</sub>NO<sub>3</sub>S.

Calc: C, 64.33; H, 6.03; N, 4.42.

25

Found: C, 64.39; H, 5.98; N, 4.46.

B. 3-(4-tert-butylbenzoyl)amino-2-thiophenecarboxylic acid

- 40 -

A solution of methyl 3-(4-tert-butylbenzoyl)amino-2-thiophenecarboxylate (9.67 g, 30 mmol) in 1,4-dioxane (75 mL) was treated with 2 N aqueous sodium hydroxide (75 mL). After 16 h, the mixture was treated with 5 N aqueous  
5 hydrochloric acid until the pH was ~2. The mixture was poured into ethyl acetate and the layers separated. The aqueous layer was washed three times with ethyl acetate and the combined extracts were dried (magnesium sulfate), filtered, and concentrated in vacuo to yield 8.09 g (89%) of  
10 the title compound.

<sup>1</sup>H-NMR, IR

MS-FD m/e 303 (M<sup>+</sup>)

Analysis for C<sub>16</sub>H<sub>17</sub>NO<sub>3</sub>S.

Calc: C, 63.34; H, 5.65; N, 4.62.

15 Found: C, 63.56; H, 5.93; N, 4.32.

C. 2-[4-(tert-butyl)phenyl]-4-oxo-4H-thieno[3,2-d]-[1.3]oxazine

A solution of 3-(4-tert-butylbenzoyl)amino-2-  
20 thiophenecarboxylic acid (8.1 g, 27 mmol) in methylene chloride (135 mL) was treated with oxalyl chloride (11.8 mL, 135 mmol). The mixture was slowly heated to afford a homogeneous solution. After 2 h, the mixture was concentrated in vacuo and the residue dissolved in methylene  
25 chloride (135 mL) and treated with pyridine (2.2 mL). After 1 hr, the mixture was concentrated in vacuo and the residue partitioned between ethyl acetate and water. The organic layer was washed four times with water, once with saturated sodium chloride solution, dried (magnesium sulfate), and  
30 filtered. Concentration and purification of the residue by flash chromatography (silica gel, hexanes/ethyl acetate) yielded 7.44 g (96%) of the title compound.

- 41 -

<sup>1</sup>H-NMR, IRMS-FD m/e 285 (M<sup>+</sup>)Analysis for C<sub>16</sub>H<sub>15</sub>NO<sub>2</sub>S.

Calc: C, 67.34; H, 5.30; N, 4.91.

5 Found: C, 67.51; H, 5.56; N, 4.76.

D. 3-(4-tert-butylbenzoyl)amino-N-(4-methoxyphenyl)-2-thiophenecarboxamide

A solution of 2-[4-(tert-butyl)phenyl]-4-oxo-4H-thieno[3,2-d][1.3]oxazine (60 mg, 0.21 mmol) and p-anisidine (26 mg, 0.21 mmol) in toluene (1 mL) was treated with p-toluenesulfonic acid (4 mg) and the resulting mixture heated at reflux for 30 h. The mixture was concentrated in vacuo and the residue was purified by flash chromatography (silica gel, hexanes/ethyl acetate) to yield 35 mg (41%) of the title compound.


<sup>1</sup>H-NMRMS-FD m/e 408 (M<sup>+</sup>)Analysis for C<sub>23</sub>H<sub>24</sub>N<sub>2</sub>O<sub>3</sub>S.

20 Calc: C, 67.62; H, 5.92; N, 6.86.

Found: C, 67.79; H, 5.84; N, 6.77.

### Example 2

Preparation of N<sup>3</sup>-(4-tert-butylbenzoyl)-N<sup>2</sup>-(4-methoxybenzoyl)-2,3-pyridinediamine.



- 42 -

A.  $N^3$ -(tert-butoxycarbonyl)- $N^2$ -(4-methoxybenzoyl)-2,3-pyridinediamine

A solution of  $N^3$ -(tert-butoxycarbonyl)-2,3-pyridine-  
5 diamine (446 mg, 2.13 mmol) in tetrahydrofuran (7 mL) was  
treated with potassium hexamethyldisilazide (894 mg, 4.48  
mmol). After 0.1 hr, the mixture was treated with *p*-anisoyl  
chloride (0.365 mL, 2.13 mmol). After 0.5 hr, the mixture  
was poured into aqueous ammonium chloride solution and ethyl  
10 acetate. The organic layer was washed three times with  
water, once with saturated sodium chloride solution, dried  
(magnesium sulfate), and filtered. Concentration in vacuo  
and purification of the residue by flash chromatography  
(silica gel, hexanes/ethyl acetate) yielded 250 mg (34%) of  
15 the title compound.

$^1\text{H-NMR}$

B.  $N^2$ -(4-methoxybenzoyl)-2,3-pyridinediamine

A solution of  $N^3$ -(tert-butoxycarbonyl)- $N^2$ -(4-methoxy-  
20 benzoyl)-2,3-pyridinediamine (350 mg, 1.02 mmol) in acetic  
acid (2 mL) at 0 °C was treated with boron trifluoride  
etherate (0.50 mL, 4.1 mmol). After 2 h, the mixture was  
poured into aqueous sodium bicarbonate and ethyl acetate.  
The aqueous layer was extracted three times with ethyl  
25 acetate and the combined extracts were dried (magnesium  
sulfate), filtered, and concentrated in vacuo. Purification  
of the residue by chromatography (silica gel, ethyl  
acetate/methylene chloride) yielded 122 mg (49%) of the  
title compound.

30  $^1\text{H-NMR}$

- 43 -

C.  $N^3$ -(4-tert-butylbenzoyl)- $N^2$ -(4-methoxybenzoyl)-2,3-pyridinediamine

A solution of  $N^2$ -(4-methoxybenzoyl)-2,3-pyridinediamine (92 mg, 0.38 mmol) in tetrahydrofuran (1 mL) was treated with potassium hexamethyldisilazide (160 mg, 0.80 mmol). After 0.25 hr, the mixture was treated with 4-tert-butylbenzoyl chloride. After 0.75 hr, the mixture was poured into a mixture of aqueous ammonium chloride solution and ethyl acetate. The organic layer was washed twice with water, once with saturated sodium chloride solution, dried (magnesium sulfate), and filtered. Concentration in vacuo and purification of the residue by flash chromatography (silica gel, methylene chloride/ethyl acetate) followed by recrystallization from ethyl acetate/hexanes yielded 31 mg (20%) of the title compound.

$^1\text{H-NMR}$

MS-FD m/e 403 ( $M^+$ )


Analysis for  $\text{C}_{24}\text{H}_{25}\text{N}_3\text{O}_3$ .

Calc: C, 71.44; H, 6.24; N, 10.41.

Found: C, 71.28; H, 6.33; N, 10.52.

Example 3

Preparation of  $N^4$ -(4-tert-butylbenzoyl)- $N^3$ -(4-methoxybenzoyl)-3,4-pyridinediamine.



A.  $N^4$ -(tert-butyloxycarbonyl)-3,4-pyridinediamine

- 44 -

A solution of 3,4-pyridinediamine (930 mg, 8.52 mmol) in tetrahydrofuran (20 mL) was treated with water (20 mL) and potassium carbonate (2.35 g, 17.0 mmol) followed by di-tert-butyl dicarbonate (1.86 g, 8.52 mmol). After 0.75 hr, the mixture was poured into a mixture of ethyl acetate and water. The aqueous layer was saturated with sodium chloride and extracted several times with ethyl acetate. The combined organic extracts were washed with saturated sodium chloride solution, dried (magnesium sulfate), and filtered. Concentration in vacuo and recrystallization from hexanes/ethyl acetate yielded 950 mg (53%) of the title compound.

<sup>1</sup>H-NMR, IR

MS-FD m/e 209 (M<sup>+</sup>)

15 Analysis for C<sub>10</sub>H<sub>15</sub>N<sub>3</sub>O<sub>2</sub>.

Calc: C, 57.40; H, 7.23; N, 20.08.

Found: C, 57.36; H, 7.19; N, 20.29.

20 B. N<sup>4</sup>-(tert-butyloxycarbonyl)-N<sup>3</sup>-(4-methoxybenzoyl)-3,4-pyridinediamine

A solution of N<sup>4</sup>-(tert-butyloxycarbonyl)-3,4-pyridinediamine (500 mg, 2.40 mmol) and pyridine (0.213 mL, 2.63 mmol) in methylene chloride (12 mL) was treated with p-anisoyl chloride (408 mg, 2.40 mmol). After 0.75 hr, the mixture was poured into a mixture of ethyl acetate and 1 N aqueous sodium hydroxide. The organic layer was washed with saturated sodium chloride, dried (magnesium sulfate), and filtered. Concentration in vacuo and recrystallization of the residue, followed by flash chromatography (silica gel, hexanes/ethyl acetate) yielded 500 mg (60%) of the title compound.

- 45 -

<sup>1</sup>H-NMR, IRMS-FD m/e 343 (M<sup>+</sup>)Analysis for C<sub>18</sub>H<sub>21</sub>N<sub>3</sub>O<sub>4</sub>.

Calc: C, 62.96; H, 6.16; N, 12.24.

5 Found: C, 62.18; H, 6.06; N, 11.68.

C. N<sup>3</sup>-(4-methoxybenzoyl)-3,4-pyridinediamine

A solution of N<sup>4</sup>-(tert-butyloxycarbonyl)-N<sup>3</sup>-(4-methoxybenzoyl)-3,4-pyridinediamine (600 mg, 1.75 mmol) in  
10 methylene chloride (8 mL) was treated with trifluoroacetic acid (1.35 mL, 17.5 mmol). After 6.5 h, the mixture was concentrated and the residue dissolved in water and treated with 5 N aqueous sodium hydroxide. The resulting precipitate was collected by filtration yielding 325 mg  
15 (76%) of the title compound.

<sup>1</sup>H-NMR, IRMS-FD m/e 243 (M<sup>+</sup>)Analysis for C<sub>13</sub>H<sub>13</sub>N<sub>3</sub>O<sub>2</sub>.

Calc: C, 64.19; H, 5.39; N, 17.27.

20 Found: C, 63.92; H, 5.28; N, 17.15.

D. N<sup>4</sup>-(4-tert-butylbenzoyl)-N<sup>3</sup>-(4-methoxybenzoyl)-3,4-pyridinediamine

Using 4-tert-butylbenzoyl chloride and a similar  
25 procedure to that described for Example 2, Part C, N<sup>3</sup>-(4-methoxybenzoyl)-3,4-pyridinediamine (300 mg, 1.23 mmol) yielded 243 mg (49%) of the title compound.

<sup>1</sup>H-NMR, IRMS-FD m/e 403 (M<sup>+</sup>)30 Analysis for C<sub>24</sub>H<sub>25</sub>N<sub>3</sub>O<sub>3</sub>.


Calc: C, 71.44; H, 6.25; N, 10.41.

- 46 -

Found: C, 71.28; H, 6.16; N, 10.28.

## Example 4

Preparation of N<sup>3</sup>-(4-tert-butylbenzoyl)-N<sup>4</sup>-(4-methoxy-  
5 benzoyl)-3,4-pyridinediamine.



A. N<sup>3</sup>-(4-tert-butylbenzoyl)-N<sup>4</sup>-(tert-butyloxycarbonyl)-  
10 3,4-pyridinediamine

Using a similar procedure to that described for  
Example 3, Part B, N<sup>4</sup>-(tert-butyloxycarbonyl)-3,4-  
pyridinediamine (450 mg, 2.15 mmol) yielded 516 mg (65%) of  
the title compound.

15 <sup>1</sup>H-NMR, IR

MS-FD m/e 369 (M<sup>+</sup>)

Analysis for C<sub>21</sub>H<sub>27</sub>N<sub>3</sub>O<sub>3</sub>.

Calc: C, 68.27; H, 7.37; N, 11.37.

Found: C, 68.46; H, 7.38; N, 11.19.

20

B. N<sup>3</sup>-(4-tert-butylbenzoyl)-3,4-pyridinediamine

Using a similar procedure to that described for  
Example 3, Part C, N<sup>3</sup>-(4-tert-butylbenzoyl)-N<sup>4</sup>-(tert-  
butyloxycarbonyl)-3,4-pyridinediamine (516 mg, 1.40 mmol)  
25 yielded 324 mg (86%) of the title compound.

<sup>1</sup>H-NMR, IR

MS-FD m/e 269 (M<sup>+</sup>)



- 47 -

Analysis for  $C_{16}H_{19}N_3O$ .

Calc: C, 71.35; H, 7.11; N, 15.60.

Found: C, 71.01; H, 7.06; N, 14.90.

5 C.  $N^3$ -(4-*tert*-butylbenzoyl)- $N^4$ -(4-methoxybenzoyl)-3,4-pyridinediamine

A solution of  $N^3$ -(4-*tert*-butylbenzoyl)-3,4-pyridinediamine (400 mg, 1.49 mmol), pyridine (0.264 mL), and *p*-anisoyl chloride (0.254 mL, 1.49 mmol) in toluene  
10 (12 mL) was heated at reflux for 3 h. The mixture was filtered and the filtrate poured into a mixture of ethyl acetate and water. The organic layer was washed three times with water, once with saturated sodium chloride solution, dried, and filtered. Concentration in vacuo and  
15 purification of the residue by recrystallization (methanol/ethyl acetate/hexanes) followed by flash chromatography (silica gel, ethyl acetate/methylene chloride) yielded 75 mg (13%) of the title compound.  
 $^1H$ -NMR, IR

20 MS-FD m/e 403 (M+)

Analysis for  $C_{24}H_{25}N_3O_3$ .

Calc: C, 71.44; H, 6.24; N, 10.41.


Found: C, 69.90; H, 5.95; N, 10.25.

25

Example 5

Preparation of  $N^2$ -(4-*tert*-butylbenzoyl)- $N^3$ -(4-methoxybenzoyl)-2,3-pyridinediamine.

- 48 -

A.  $N^2$ -(4-tert-butylbenzoyl)-2,3-pyridinediamine

Using a similar procedure to that described for  
5 Example 3, Part B,  $N^3$ -(tert-butyloxycarbonyl)-2,3-pyridinediamine (1.00 g, 4.78 mmol) yielded  $N^2$ -(4-tert-butylbenzoyl)- $N^3$ -(tert-butyloxycarbonyl)-2,3-pyridinediamine (894 mg). Using a similar procedure to that described for  
10 Example 2, Part B, this crude material yielded 400 mg (31%) of the title compound.

 $^1\text{H-NMR}$ B.  $N^2$ -(4-tert-butylbenzoyl)- $N^3$ -(4-methoxybenzoyl)-2,3-pyridinediamine

15 Using a similar procedure to that described for Example 3, Part C,  $N^2$ -(4-tert-butylbenzoyl)-2,3-pyridinediamine (80 mg, 0.30 mmol) yielded 28 mg (23%) of the title compound.

 $^1\text{H-NMR}$ 

20 MS-FD m/e 403 (M+)

Analysis for  $\text{C}_{24}\text{H}_{25}\text{N}_3\text{O}_3$ .

Calc: C, 71.44; H, 6.25; N, 10.41.


Found: C, 71.51; H, 6.28; N, 10.31.

25

## Example 6

Preparation of  $N^4$ -(4-methoxybenzoyl)- $N^3$ -[4-(4-pyridyl)-benz yl]-3,4-pyridinediamine.

- 49 -



## A. sodium 4-(4-pyridyl)benzoate

5 A solution of 4-chloropyridine hydrochloride (3.00 g, 20.0 mmol), 4-carboxybenzeneboronic acid (4.97 g, 30.0 mmol), 1 M aqueous sodium carbonate solution (50 mL), 1,4-bis(diphenylphosphino)butane palladium(II) dichloride (300 mg, 0.70 mmol), and ethanol (10 mL) in toluene (40 mL) was heated at reflux for 16 h. The mixture was diluted with methanol and filtered through diatomaceous earth. The filtrate was concentrated in vacuo and the pH was adjusted to 14 by the addition of 1 N aqueous sodium hydroxide. After heating the filtrate to boiling, the insoluble material was removed by filtration, and the resulting filtrate was allowed to cool to room temperature. The resulting precipitate was collected by filtration to yield 1.83 g (41%) of the title compound.

<sup>1</sup>H-NMRB. N<sup>4</sup>-(tert-butyloxycarbonyl)-N<sup>3</sup>-[4-(4-pyridyl)benzoyl]-3,4-pyridinediamine

20 A suspension of sodium 4-(4-pyridyl)benzoate (425 mg, 1.92 mmol) in methylene chloride was treated with oxalyl chloride (0.840 mL, 9.60 mmol), followed by dimethylformamide (0.01 mL). After 0.75 h, the mixture was concentrated in vacuo. The residue was then dissolved in

- 50 -

methylene chloride and added dropwise to a solution of N<sup>4</sup>-  
(tert-butyloxycarbonyl)-3,4-pyridinediamine (400 mg, 1.92  
mmol) and pyridine (0.31 mL) in methylene chloride (2 mL)  
and tetrahydrofuran (1 mL). After 16 h, the mixture was  
5 poured into ethyl acetate and 1 N aqueous sodium hydroxide.  
The organic layer was washed once with 1 N aqueous sodium  
hydroxide, once with saturated sodium chloride solution,  
dried (potassium carbonate), and filtered. The residue was  
purified by flash chromatography (silica gel, ethyl  
10 acetate/hexanes) to yield 75 mg (10%) of the title compound.

<sup>1</sup>H-NMR, IRMS-FD m/e 390 (M<sup>+</sup>)Analysis for C<sub>22</sub>H<sub>22</sub>N<sub>4</sub>O<sub>3</sub>.

Calc: C, 67.68; H, 5.68; N, 14.35.

15 Found: C, 66.95; H, 6.03; N, 13.67.

C. N<sup>3</sup>-[4-(4-pyridyl)benzoyl]-3,4-pyridinediamine

Using a similar procedure to that described for  
Example 2, Part B, N<sup>4</sup>-(tert-butyloxycarbonyl)-N<sup>3</sup>-[4-  
20 (4-pyridyl)benzoyl]-3,4-pyridinediamine (95 mg, 0.23 mmol)  
yielded 55 mg (82%) of the title compound.

<sup>1</sup>H-NMR

D. N<sup>4</sup>-(4-methoxybenzoyl)-N<sup>3</sup>-[4-(4-pyridyl)benzoyl]-3,4-  
25 pyridinediamine

Using 4-methoxybenzoyl chloride and a similar procedure  
to that described for Example 2, Part C, N<sup>3</sup>-[4-(4-pyridyl)-  
benzoyl]-3,4-pyridinediamine (55 mg, 0.19 mmol) yielded 3.2  
mg (4%) of the title compound.

30 <sup>1</sup>H-NMRMS-FD m/e 424 (M<sup>+</sup>)

- 51 -


## Examples 7-9

The following procedure was use in Examples 7-9:

5 To a small glass vial with a Teflon lined cap was added a 1,2-diamino aromatic compound (about 0.25 mmol) in tetrahydrofuran (3 mL), followed by poly(4-vinylpyridine) (250 mg, 1 mmol) and *p*-anisoyl chloride (0.625 mmol). After  
10 agitating this mixture for 24 h on a platform shaker, aminomethylated polystyrene (1 g, 1 mmol) was added and agitation continued for another 8 h. The solution was filtered and concen-  
15 trated in vacuo, and the residue triturated with diethyl ether. The resulting solid was filtered and dried in vacuo to give approximately 50 mg of the title compound.

## Example 7

5-Bromo-6-methyl-*N*<sup>2</sup>,*N*<sup>3</sup>-bis(4-methoxybenzoyl)-2,3-pyridine-  
20 diamine.




MS-FD m/e 471 (M<sup>+</sup>).

- 52 -

## Example 8

**N<sup>3</sup>,N<sup>4</sup>-bis(4-Methoxybenzoyl)-3,4-pyridinediamine.**




5

MS-FD m/e 377 (M<sup>+</sup>).

## Example 9

**N<sup>2</sup>,N<sup>3</sup>-bis(4-Methoxybenzoyl)-2,3-pyridinediamine.**




10

MS-FD m/e 377 (M<sup>+</sup>).

## Example 10

15 **Preparation of 3-(4-Methoxybenzoyl)amino-N-(4-methoxyphenyl)-2-indolecarboxamide.**

- 53 -



A. 3-(4-methoxybenzoyl)amino-2-indolecarboxylic acid ethyl ester

5 To a solution of 3-amino-2-indolecarboxylic acid ethyl ester (500 mg, 2.45 mmol) and triethylamine (272 mg, 2.70 mmol) in methylene chloride (5 mL) was added anisoyl chloride (418 mg, 2.45 mmol). The resulting mixture was stirred for 18 h at room temperature. The reaction mixture  
10 was diluted with ethyl acetate and washed with water. The organic phase was washed with 1 N aqueous hydrochloric acid, dried (sodium sulfate), filtered, and concentrated in vacuo to provide a yellow solid. Recrystallization from  
15 hexane/ethyl acetate provided 780 mg (94%) of the title compound as a yellow solid.

<sup>1</sup>H-NMR, IR

MS-FD m/e 338 (M<sup>+</sup>)

Analysis for C<sub>19</sub>H<sub>18</sub>N<sub>2</sub>O<sub>4</sub>.

Calc: C, 67.44; H, 5.36; N, 8.28.

20 Found: C, 67.46; H, 5.35; N, 8.16.

B. 3-(4-methoxybenzoyl)amino-2-indolecarboxylic acid

To a solution of 3-(4-methoxybenzoyl)amino-2-indolecarboxylic acid ethyl ester (720 mg, 2.13 mmol) in

- 54 -

tetrahydrofuran (7 mL) was added 5 N aqueous sodium hydroxide (2 mL). The resulting mixture was stirred for 10 h at room temperature. An additional portion of 5 N aqueous sodium hydroxide (5 mL) was added and the mixture  
5 heated at 60 °C for 5 h. The mixture was cooled to room temperature, stirred for 10 h, diluted with water, and extracted with diethyl ether. The aqueous layer was acidified with concentrated hydrochloric acid and extracted with three fresh portions of ethyl acetate. The combined  
10 ethyl acetate fractions were dried (sodium sulfate), filtered, and concentrated in vacuo to provide 450 mg (68%) of the title compound as a yellow solid.

<sup>1</sup>H-NMR, IR

MS-FD m/e 310 (M<sup>+</sup>)

15 Analysis for C<sub>17</sub>H<sub>14</sub>N<sub>2</sub>O<sub>4</sub>.

Calc: C, 65.80; H, 4.55; N, 9.03.

Found: C, 64.70; H, 4.66; N, 8.59.

C. 3-(4-methoxybenzoyl)amino-N-(4-methoxyphenyl)-2-  
20 indolecarboxamide

To a solution p-anisidine (79 mg, 0.645 mmol) in methylene chloride (5 mL) was added 3-(4-methoxybenzoyl)-amino-2-indolecarboxylic acid (200 mg, 0.645 mmol),  
1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride  
25 (247 mg, 1.29 mmol), and 4-dimethylaminopyridine (8.0 mg, 0.065 mmol). The resulting solution was stirred at room temperature for 6 h. The resulting precipitate was collected via vacuum filtration to provide 38 mg (14%) of the title compound as a white solid.

30 <sup>1</sup>H-NMR, IR

MS-FD m/e 415 (M<sup>+</sup>)



- 55 -

Analysis for  $C_{24}H_{21}N_3O_4$ .


Calc: C, 69.39; H, 5.10; N, 10.11.

Found: C, 68.68; H, 4.96; N, 10.15.

5

## Example 11

Preparation of  $N^4$ -[(4-Dimethylamino)benzoyl]- $N^3$ -(4-methoxybenzoyl)-3,4-pyridinediamine.



10 A. 4-(Dimethylamino)benzoyl chloride

A solution of 4-(dimethylamino)benzoic acid and thionyl chloride in methylene chloride was refluxed 4 h. Volatile solvents were removed in vacuo to yield 1.10 g of 4-(dimethylamino)benzoyl chloride. This material was used in subsequent reactions without purification.

B.  $N^4$ -[(4-Dimethylamino)benzoyl]- $N^3$ -(4-methoxybenzoyl)-3,4-pyridinediamine.

To a solution of  $N^3$ -(4-methoxybenzoyl)-3,4-pyridine-diamine (193 mg, 0.79 mmol) and 4-(dimethylamino)benzoyl chloride (183 mg, 1.00 mmol) in 5 mL methylene chloride was added 0.5 mL pyridine and a catalytic amount of 4-dimethylaminopyridine. The mixture was stirred 16 h at ambient temperature under nitrogen then partitioned between

- 56 -

methylene chloride and saturated sodium hydrogen carbonate solution. The organic portion was dried over magnesium sulfate and concentrated in vacuo. The residue was dissolved in ethyl acetate and hexane added until cloudy.

- 5 The mixture was sonicated inducing crystallization. The solid was collected by filtration and dried under vacuum to yield 306 mg (99%) of the title compound.

MS, Ion spray, m/e: 391(p+1).


Analysis for  $C_{22}H_{22}N_4O_3$ .

- 10        Calc.:    C, 67.58; H, 5.68; N, 14.35.  
         Found:    C, 67.19; H, 6.01; N, 13.79.

- 57 -

What is claimed is:

1. A method of inhibiting factor Xa comprising using an effective amount of a factor Xa inhibiting compound of formula I



wherein

- 10  $A^3$ ,  $A^4$ ,  $A^5$  and  $A^6$ , together with the two carbons to which they are attached, complete a substituted heteroaromatic ring in which
- (a) one of  $A^3$ ,  $A^4$ ,  $A^5$  and  $A^6$  is N, and each of the others is  $CR^3$ ,  $CR^4$ ,  $CR^5$  or  $CR^6$ , respectively;
- 15 (b) two adjacent residues of  $A^3$ ,  $A^4$ ,  $A^5$  and  $A^6$  together form S, and each of the others is  $CR^3$ ,  $CR^4$ ,  $CR^5$  or  $CR^6$ , respectively;
- (c) two non-adjacent residues of  $A^3$ ,  $A^4$ ,  $A^5$  and  $A^6$  are each N, and each of the others is  $CR^3$ ,  $CR^4$ ,  $CR^5$  or  $CR^6$ ,
- 20 respectively; or
- (d)  $A^3$  and  $A^4$  together form a fused benz ring, and  $A^5$  and  $A^6$  together form -NH-;

wherein


- each of  $R^3$ ,  $R^4$ ,  $R^5$  and  $R^6$  is hydrogen, or one or two of
- 25  $R^3$ ,  $R^4$ ,  $R^5$  and  $R^6$  is independently chloro, bromo or methyl and the others are hydrogen;

$L^1$  is -NH-CO- or -CO-NH- such that  $-L^1-Q^1$  is -NH-CO- $Q^1$  or -CO-NH- $Q^1$ ;

- $Q^1$  is phenyl, 2-furanyl, 2-thienyl, 4-thiazolyl, 2-pyridyl, 2-naphthyl, 1,2-dihydrobenzofuran-5-yl, 1,2-dihydrobenzofuran-6-yl or 1,2-benzisoxazol-6-yl in which the phenyl may bear a 2-fluoro substituent or may bear one,
- 30


- 58 -

- two or three substituents at the 3-, 4- or 5-position(s) independently selected from halo, cyano, carbamoyl, aminomethyl, methyl, methoxy, difluoromethoxy, hydroxymethyl, formyl, vinyl, amino, hydroxy and
- 5 3,4-methylenedioxy, the 2-furanyl or 2-thienyl may bear a chloro or methyl substituent at the 5-position, the 4-thiazolyl may bear an amino substituent at the 2-position, the 2-pyridyl may bear an amino substituent at the 6-position, and the 1,2-benzisoxazol-6-yl may bear a chloro
- 10 or methyl substituent at the 3-position; or -CO-Q<sup>1</sup> is cyclopentenylcarbonyl or cyclohexenylcarbonyl;
- R<sup>2</sup> is -L<sup>2A</sup>-Q<sup>2A</sup>, -L<sup>2B</sup>-Q<sup>2B</sup>, -L<sup>2C</sup>-Q<sup>2C</sup> or -L<sup>2D</sup>-Q<sup>2D</sup> wherein L<sup>2A</sup> is a direct bond; and Q<sup>2A</sup> is



- in which D is carbonyl or -CHR<sup>k</sup>- in which R<sup>k</sup> is hydrogen, hydroxy, (1-6C)alkoxy or -CH<sub>2</sub>-R<sup>j</sup> in which R<sup>j</sup> is carboxy, [(1-4C)alkoxy]carbonyl or carbamoyl which may bear one or
- 20 two (1-2C)alkyl substituents on the nitrogen; and one of R<sup>m</sup> and R<sup>n</sup> is hydrogen and the other is amino, bromo, (1-4C)alkyl or (1-4C)alkoxy, or R<sup>m</sup> and R<sup>n</sup> together form a benz ring;

- L<sup>2B</sup> is -NH-CO-, -O-CO-, -CH<sub>2</sub>-O- or -O-CH<sub>2</sub>- such that
- 25 -L<sup>2B</sup>-Q<sup>2B</sup> is -NH-CO-Q<sup>2B</sup>, -O-CO-Q<sup>2B</sup>, -CH<sub>2</sub>-O-Q<sup>2B</sup> or -O-CH<sub>2</sub>-Q<sup>2B</sup>; and

Q<sup>2B</sup> is

- 59 -

- in which  $R^O$  is hydrogen, halo, (1-6C)alkyl, (1-4C)alkoxy, benzyloxy or (1-4C)alkylthio; and  $R^P$  is 1-hydroxyethyl, 1-hydroxy-1-methylethyl, 1-methoxy-1-methylethyl,
- 5 4-piperidinyl, 4-pyridinyl, dimethylaminosulfonyl or  $-J-R^Q$  in which J is a single bond, methylene, carbonyl, oxo,  $-S(O)_q-$  (wherein q is 0, 1 or 2), or  $-NR^I-$  (wherein  $R^I$  is hydrogen or methyl); and  $R^Q$  is (1-6C)alkyl, phenyl, 3-pyridyl or 4-pyridyl;
- 10  $L^{2C}$  is  $-NR^V-CO-X-$ ,  $-NR^V-CS-Y-$ ,  $-CH_2-CO-NR^W-CH_2-$ ,  $-O-CO-$ ,  $-O-CH_2-$ ,  $-S-CH_2-$  or  $-CH_2-NR^X-CH_2-$  such that  $-L^{2C}-Q^{2C}$  is  $-NR^V-CO-X-Q^{2C}$ ,  $-NR^V-CS-Y-Q^{2C}$ ,  $-CH_2-CO-NR^W-CH_2-Q^{2C}$ ,  $-O-CO-Q^{2C}$ ,  $-O-CH_2-Q^{2C}$ ,  $-S-CH_2-Q^{2C}$  or  $-CH_2-NR^X-CH_2-Q^{2C}$  in which X is  $-(CH_2)_x-$  (wherein x is 0, 1 or 2),  $-NR^W-$ ,
- 15  $-NR^W-CH_2-$ ,  $-O-$ ,  $-O-CH_2-$  or  $-S-CH_2-$ ; Y is  $-NR^W-CH_2-$  or  $-O-CH_2-$ ; each of  $R^V$  and  $R^W$  is independently hydrogen, benzyl or (1-6C)alkyl which is not branched at the  $\alpha$ -position; and  $R^X$  is hydrogen, benzyloxycarbonyl or [(1-4C)alkoxy]carbonyl; and
- 20  $Q^{2C}$  is 1-(4-pyridyl)piperidin-4-yl, 1-(4-pyridyl)piperidin-3-yl or 1-(4-pyridyl)pyrrolidin-3-yl in which the pyridyl may bear a substituent at its 2-position selected from cyano, aminomethyl, carboxy, hydroxymethyl and (1-2C)alkyl;
- 25  $L^{2D}$  is  $-NH-CO-$  such that  $-L^{2D}-Q^{2D}$  is  $-NH-CO-Q^{2D}$ ; and  $Q^{2D}$  is selected from 4-(4-pyridinyl)benzyloxy, 9-oxo-9H-fluoren-3-yl, benzo[b]thiophen-2-yl (which may bear a chloro, methyl or methoxy substituent), benzofuran-2-yl (which may bear a chloro, methyl or methoxy substituent),
- 30 4-(4-morpholinyl)-4-oxobutyl, and 4-piperidinyl or 3,4-didehydropiperidin-4-yl (either one bearing a substituent at the 1-position selected from methylsulfonyl, phenylsulfonyl, (1-5C)alkyl, (4-7C)cycloalkyl, tetrahydropyran-4-yl, 4-thiacyclohexyl and  $-CH_2-R^Z$  in which  $R^Z$  is

- 60 -

isopropyl, cyclopropyl, phenyl, furyl, thienyl, 2-thiazolyl, or pyridyl in which the phenyl may bear one or two substituents independently selected from halo, cyano, hydroxy, methoxy, acetoxy, benzyloxy, amino, acetylamino, 5 nitro and 3,4-methylenedioxy, and the thienyl or furyl may bear a methyl or nitro substituent);

or a prodrug of the compound of formula I;

or a pharmaceutically acceptable salt of the compound of formula I or prodrug thereof.

10

2. The method of claim 1 in which the factor Xa inhibiting compound of formula I is one wherein

A<sup>3</sup>, A<sup>4</sup>, A<sup>5</sup> and A<sup>6</sup>, together with the two carbons to which they are attached, complete a substituted 15 heteroaromatic ring in which

(a) one of A<sup>3</sup>, A<sup>4</sup>, A<sup>5</sup> and A<sup>6</sup> is N, and each of the others is CR<sup>3</sup>, CR<sup>4</sup>, CR<sup>5</sup> or CR<sup>6</sup>, respectively;

(b) two adjacent residues of A<sup>3</sup>, A<sup>4</sup>, A<sup>5</sup> and A<sup>6</sup> together form S, and each of the others is CR<sup>3</sup>, CR<sup>4</sup>, CR<sup>5</sup> or CR<sup>6</sup>, 20 respectively;

(c) two non-adjacent residues of A<sup>3</sup>, A<sup>4</sup>, A<sup>5</sup> and A<sup>6</sup> are each N, and each of the others is CR<sup>3</sup>, CR<sup>4</sup>, CR<sup>5</sup> or CR<sup>6</sup>, respectively; or

(d) A<sup>3</sup> and A<sup>4</sup> together form a fused benz ring, and A<sup>5</sup> and 25 A<sup>6</sup> together form -NH-;

wherein

each of R<sup>3</sup>, R<sup>4</sup>, R<sup>5</sup> and R<sup>6</sup> is hydrogen, or one or two of R<sup>3</sup>, R<sup>4</sup>, R<sup>5</sup> and R<sup>6</sup> is independently chloro, bromo or methyl and the others are hydrogen;


30 L<sup>1</sup> is -NH-CO- or -CO-NH- such that -L<sup>1</sup>-Q<sup>1</sup> is -NH-CO-Q<sup>1</sup> or -CO-NH-Q<sup>1</sup>;

Q<sup>1</sup> is phenyl, 2-thienyl, 4-thiazolyl, 2-pyridyl, 2-naphthyl or 1,2-benzisoxazol-6-yl in which the phenyl may bear one, two or three substituents at the 3-, 4- or

- 61 -

5-position(s) independently selected from halo, cyano, carbamoyl, aminomethyl, methyl, methoxy, hydroxymethyl, formyl, vinyl, amino, hydroxy and 3,4-methylenedioxy, the 2-thienyl may bear a chloro or methyl substituent at the 5-position, the 4-thiazolyl may bear an amino substituent at the 2-position, the 2-pyridyl may bear an amino substituent at the 6-position, and the 1,2-benzisoxazol-6-yl may bear a chloro or methyl substituent at the 3-position;


$R^2$  is  $-L^{2A}-Q^{2A}$ ,  $-L^{2B}-Q^{2B}$ ,  $-L^{2C}-Q^{2C}$  or  $-L^{2D}-Q^{2D}$  wherein  
 10  $L^{2A}$  is a direct bond; and  
 $Q^{2A}$  is



in which D is carbonyl or  $-CHR^k-$  in which  $R^k$  is hydrogen, hydroxy, (1-6C)alkoxy or  $-CH_2-R^j$  in which  $R^j$  is carboxy, [(1-4C)alkoxy]carbonyl or carbamoyl which may bear one or two (1-2C)alkyl substituents on the nitrogen; and one of  $R^m$  and  $R^n$  is hydrogen and the other is amino, bromo, (1-4C)alkyl or (1-4C)alkoxy, or  $R^m$  and  $R^n$  together form a benz ring;

$L^{2B}$  is  $-NH-CO-$ ,  $-O-CO-$ ,  $-CH_2-O-$  or  $-O-CH_2-$  such that  $-L^{2B}-Q^{2B}$  is  $-NH-CO-Q^{2B}$ ,  $-O-CO-Q^{2B}$ ,  $-CH_2-O-Q^{2B}$  or  $-O-CH_2-Q^{2B}$ ; and

$Q^{2B}$  is



25

in which  $R^o$  is hydrogen, halo, (1-6C)alkyl, (1-4C)alkoxy, benzyloxy or (1-4C)alkylthio; and  $R^p$  is 1-hydroxyethyl, 1-hydroxy-1-methylethyl, 1-methoxy-1-methylethyl,

- 62 -

- 4-piperidinyl, 4-pyridinyl, dimethylaminosulfonyl or  $-J-R^Q$  in which J is a single bond, methylene, carbonyl, oxo,  $-S(O)_q-$  (wherein q is 0, 1 or 2), or  $-NR^T-$  (wherein  $R^T$  is hydrogen or methyl); and  $R^Q$  is (1-6C)alkyl, phenyl, 3-pyridyl or 4-pyridyl;
- 5  $L^{2C}$  is  $-NR^V-CO-X-$ ,  $-NR^V-CS-Y-$ ,  $-CH_2-CO-NR^W-CH_2-$ ,  $-O-CO-$ ,  $-O-CH_2-$ ,  $-S-CH_2-$  or  $-CH_2-NR^X-CH_2-$  such that  $-L^{2C}-Q^{2C}$  is  $-NR^V-CO-X-Q^{2C}$ ,  $-NR^V-CS-Y-Q^{2C}$ ,  $-CH_2-CO-NR^W-CH_2-Q^{2C}$ ,  $-O-CO-Q^{2C}$ ,  $-O-CH_2-Q^{2C}$ ,  $-S-CH_2-Q^{2C}$  or  $-CH_2-NR^X-CH_2-Q^{2C}$  in
- 10 which X is  $-(CH_2)_x-$  (wherein x is 0, 1 or 2),  $-NR^W-CH_2-$ ,  $-O-CH_2-$  or  $-S-CH_2-$ ; Y is  $-NR^W-CH_2-$  or  $-O-CH_2-$ ; each of  $R^V$  and  $R^W$  is independently hydrogen, benzyl or (1-6C)alkyl which is not branched at the  $\alpha$ -position; and  $R^X$  is hydrogen, benzyloxycarbonyl or [(1-4C)alkoxy]carbonyl; and
- 15  $Q^{2C}$  is 1-(4-pyridyl)piperidin-4-yl in which the pyridyl may bear a substituent at its 2-position selected from cyano, aminomethyl, carboxy, hydroxymethyl and (1-2C)alkyl;
- $L^{2D}$  is  $-NH-CO-$  such that  $-L^{2D}-Q^{2D}$  is  $-NH-CO-Q^{2D}$ ; and  $Q^{2D}$  is selected from 4-(4-pyridinyl)benzyloxy, 9-oxo-
- 20 9H-fluoren-3-yl, benzo[b]thiophen-2-yl (which may bear a chloro, methyl or methoxy substituent), benzofuran-2-yl (which may bear a chloro, methyl or methoxy substituent), 4-(4-morpholinyl)-4-oxobutyl, and 4-piperidinyl bearing a substituent at the 1-position selected from methylsulfonyl, phenylsulfonyl and  $-CH_2-R^Z$  in which  $R^Z$  is isopropyl,
- 25 cyclopropyl, phenyl, furyl, thienyl, 2-thiazolyl, or pyridyl in which the phenyl may bear one or two substituents independently selected from halo, cyano, hydroxy, methoxy, acetoxy, benzyloxy, amino, acetylamino, nitro and
- 30 3,4-methylenedioxy, and the thienyl or furyl may bear a methyl or nitro substituent;
- or a prodrug of the compound of formula I;
- or a pharmaceutically acceptable salt of the compound of formula I or prodrug thereof.



- 63 -

3. The method of Claim 1 or 2 wherein for an alkyl group or the alkyl portion of an alkyl containing group, (1-2C)alkyl is methyl or ethyl; (1-4C)alkyl is methyl, ethyl, propyl, isopropyl, butyl, isobutyl, or t-butyl; (1-6C)alkyl is methyl, ethyl, propyl, butyl, pentyl or hexyl; and halo is bromo or chloro.

4. The method of Claim 3 wherein for an alkyl group or the alkyl portion of an alkyl containing group, (1-2C)alkyl is methyl; (1-4C)alkyl is methyl, isopropyl, butyl or t-butyl; (1-6C)alkyl is methyl, butyl or hexyl; and halo is chloro.

5. The method of any of the above Claims 1-4 wherein  $R^2$  is selected from  $-L^2A-Q^2A$ ,  $-NH-CO-Q^2B$ ,  $-NR^V-CO-X-Q^2C$ ,  $-NR^V-CS-Y-Q^2C$ , and  $-NH-CO-Q^2D$ .

6. The method of any of the above Claims 1-5 wherein the compound of formula I is a pyridine in which one of  $A^3$ ,  $A^4$ ,  $A^5$  and  $A^6$  is N, and each of the others is  $CR^3$ ,  $CR^4$ ,  $CR^5$  or  $CR^6$ , respectively.

7. The method of any of the above Claims 1-6 wherein the compound of formula I is a thiophene in which the two adjacent residues  $A^5$  and  $A^6$  together form S, and  $A^3$  and  $A^4$  are  $CR^3$  and  $CR^4$ , respectively.

8. The method of any of the above Claims 1-6 wherein the compound of formula I is an indole in which the two adjacent residues  $A^5$  and  $A^6$  together form  $-NH-$ , and  $A^3$  and  $A^4$  together form a fused benz ring.

- 64 -

9. The method of Claim 6 wherein  $A^4$  is N, and each of  $R^3$ ,  $R^5$  and  $R^6$  is hydrogen.


10. The method of any of the above Claims 1-9 wherein  $Q^1$  is 4-methoxyphenyl.

11. The method of any of the above Claims 1-10 wherein  $R^2$  is (4-t-butylbenzoyl)amino, (4-methoxybenzoyl)amino, [4-(4-pyridyl)benzoyl]amino or [1-(4-pyridyl)piperidin-4-yl]methoxycarbonylamino.

12. The method of any of the above Claims 1-11 wherein  $L^1-Q^1$  is  $-NH-CO-Q^1$ .

13. The method of any of the above Claims 1-11 wherein  $L^1-Q^1$  is  $-CO-NH-Q^1$ .

14. A novel compound of formula I



wherein

- $A^3$ ,  $A^4$ ,  $A^5$  and  $A^6$ , together with the two carbons to which they are attached, complete a substituted heteroaromatic ring in which
- (a) one of  $A^3$ ,  $A^4$ ,  $A^5$  and  $A^6$  is N, and each of the others is  $CR^3$ ,  $CR^4$ ,  $CR^5$  or  $CR^6$ , respectively;
  - (b) two adjacent residues of  $A^3$ ,  $A^4$ ,  $A^5$  and  $A^6$  together form S, and each of the others is  $CR^3$ ,  $CR^4$ ,  $CR^5$  or  $CR^6$ , respectively;

- 65 -

(c) two non-adjacent residues of  $A^3$ ,  $A^4$ ,  $A^5$  and  $A^6$  are each N, and each of the others is  $CR^3$ ,  $CR^4$ ,  $CR^5$  or  $CR^6$ , respectively; or


(d)  $A^3$  and  $A^4$  together form a fused benz ring, and  $A^5$  and  $A^6$  together form -NH-;  
 5 wherein

each of  $R^3$ ,  $R^4$ ,  $R^5$  and  $R^6$  is hydrogen, or one or two of  $R^3$ ,  $R^4$ ,  $R^5$  and  $R^6$  is independently chloro, bromo or methyl and the others are hydrogen;

10  $L^1$  is -NH-CO- or -CO-NH- such that  $-L^1-Q^1$  is -NH-CO- $Q^1$  or -CO-NH- $Q^1$ ;

$Q^1$  is phenyl, 2-furanyl, 2-thienyl, 4-thiazolyl, 2-pyridyl, 2-naphthyl, 1,2-dihydrobenzofuran-5-yl, 1,2-dihydrobenzofuran-6-yl or 1,2-benzisoxazol-6-yl in which  
 15 the phenyl may bear a 2-fluoro substituent or may bear one, two or three substituents at the 3-, 4- or 5-position(s) independently selected from halo, cyano, carbamoyl, aminomethyl, methyl, methoxy, difluoromethoxy, hydroxymethyl, formyl, vinyl, amino, hydroxy and  
 20 3,4-methylenedioxy, the 2-furanyl or 2-thienyl may bear a chloro or methyl substituent at the 5-position, the 4-thiazolyl may bear an amino substituent at the 2-position, the 2-pyridyl may bear an amino substituent at the 6-position, and the 1,2-benzisoxazol-6-yl may bear a chloro  
 25 or methyl substituent at the 3-position; or -CO- $Q^1$  is cyclopentenylcarbonyl or cyclohexenylcarbonyl;


$R^2$  is - $L^{2A}$ - $Q^{2A}$ , - $L^{2B}$ - $Q^{2B}$ , - $L^{2C}$ - $Q^{2C}$  or - $L^{2D}$ - $Q^{2D}$  wherein  
 $L^{2A}$  is a direct bond; and  
 $Q^{2A}$  is



- 66 -

in which D is carbonyl or  $-\text{CHR}^k-$  in which  $\text{R}^k$  is hydrogen, hydroxy, (1-6C)alkoxy or  $-\text{CH}_2-\text{R}^j$  in which  $\text{R}^j$  is carboxy, [(1-4C)alkoxy]carbonyl or carbamoyl which may bear one or two (1-2C)alkyl substituents on the nitrogen; and one of  $\text{R}^m$  and  $\text{R}^n$  is hydrogen and the other is amino, bromo, (1-4C)alkyl or (1-4C)alkoxy, or  $\text{R}^m$  and  $\text{R}^n$  together form a benz ring;

$\text{L}^{2\text{B}}$  is  $-\text{NH}-\text{CO}-$ ,  $-\text{O}-\text{CO}-$ ,  $-\text{CH}_2-\text{O}-$  or  $-\text{O}-\text{CH}_2-$  such that  $-\text{L}^{2\text{B}}-\text{Q}^{2\text{B}}$  is  $-\text{NH}-\text{CO}-\text{Q}^{2\text{B}}$ ,  $-\text{O}-\text{CO}-\text{Q}^{2\text{B}}$ ,  $-\text{CH}_2-\text{O}-\text{Q}^{2\text{B}}$  or  $-\text{O}-\text{CH}_2-\text{Q}^{2\text{B}}$ ; and  $\text{Q}^{2\text{B}}$  is



in which  $\text{R}^{\text{O}}$  is hydrogen, halo, (1-6C)alkyl, (1-4C)alkoxy, benzyloxy or (1-4C)alkylthio; and  $\text{R}^{\text{P}}$  is 1-hydroxyethyl, 1-hydroxy-1-methylethyl, 1-methoxy-1-methylethyl, 4-piperidinyl, 4-pyridinyl, dimethylaminosulfonyl or  $-\text{J}-\text{R}^{\text{Q}}$  in which J is a single bond, methylene, carbonyl, oxo,  $-\text{S}(\text{O})_{\text{q}}-$  (wherein q is 0, 1 or 2), or  $-\text{NR}^{\text{F}}-$  (wherein  $\text{R}^{\text{F}}$  is hydrogen or methyl); and  $\text{R}^{\text{Q}}$  is (1-6C)alkyl, phenyl, 3-pyridyl or 4-pyridyl;

$\text{L}^{2\text{C}}$  is  $-\text{NR}^{\text{V}}-\text{CO}-\text{X}-$ ,  $-\text{NR}^{\text{V}}-\text{CS}-\text{Y}-$ ,  $-\text{CH}_2-\text{CO}-\text{NR}^{\text{W}}-\text{CH}_2-$ ,  $-\text{O}-\text{CO}-$ ,  $-\text{O}-\text{CH}_2-$ ,  $-\text{S}-\text{CH}_2-$  or  $-\text{CH}_2-\text{NR}^{\text{X}}-\text{CH}_2-$  such that  $-\text{L}^{2\text{C}}-\text{Q}^{2\text{C}}$  is  $-\text{NR}^{\text{V}}-\text{CO}-\text{X}-\text{Q}^{2\text{C}}$ ,  $-\text{NR}^{\text{V}}-\text{CS}-\text{Y}-\text{Q}^{2\text{C}}$ ,  $-\text{CH}_2-\text{CO}-\text{NR}^{\text{W}}-\text{CH}_2-\text{Q}^{2\text{C}}$ ,  $-\text{O}-\text{CO}-\text{Q}^{2\text{C}}$ ,  $-\text{O}-\text{CH}_2-\text{Q}^{2\text{C}}$ ,  $-\text{S}-\text{CH}_2-\text{Q}^{2\text{C}}$  or  $-\text{CH}_2-\text{NR}^{\text{X}}-\text{CH}_2-\text{Q}^{2\text{C}}$  in which X is  $-(\text{CH}_2)_x-$  (wherein x is 0, 1 or 2),  $-\text{NR}^{\text{W}}-$ ,  $-\text{NR}^{\text{W}}-\text{CH}_2-$ ,  $-\text{O}-$ ,  $-\text{O}-\text{CH}_2-$  or  $-\text{S}-\text{CH}_2-$ ; Y is  $-\text{NR}^{\text{W}}-\text{CH}_2-$  or  $-\text{O}-\text{CH}_2-$ ; each of  $\text{R}^{\text{V}}$  and  $\text{R}^{\text{W}}$  is independently hydrogen, benzyl or (1-6C)alkyl which is not branched at the  $\alpha$ -position; and  $\text{R}^{\text{X}}$  is hydrogen, benzyloxycarbonyl or [(1-4C)alkoxy]carbonyl; and

- 67 -

Q<sup>2C</sup> is 1-(4-pyridyl)piperidin-4-yl, 1-(4-pyridyl)-piperidin-3-yl or 1-(4-pyridyl)pyrrolidin-3-yl in which the pyridyl may bear a substituent at its 2-position selected from cyano, aminomethyl, carboxy, hydroxymethyl and

5 (1-2C)alkyl;

L<sup>2D</sup> is -NH-CO- such that -L<sup>2D</sup>-Q<sup>2D</sup> is -NH-CO-Q<sup>2D</sup>; and

Q<sup>2D</sup> is selected from 4-(4-pyridinyl)benzyloxy, 9-oxo-9H-fluoren-3-yl, benzo[b]thiophen-2-yl (which may bear a chloro, methyl or methoxy substituent), benzofuran-2-yl  
10 (which may bear a chloro, methyl or methoxy substituent), 4-(4-morpholinyl)-4-oxobutyl, and 4-piperidinyl or 3,4-didehydropiperidin-4-yl (either one bearing a substituent at the 1-position selected from methylsulfonyl, phenylsulfonyl, (1-5C)alkyl, (4-7C)cycloalkyl, tetrahydro-  
15 pyran-4-yl, 4-thiacyclohexyl and -CH<sub>2</sub>-R<sup>Z</sup> in which R<sup>Z</sup> is isopropyl, cyclopropyl, phenyl, furyl, thienyl, 2-thiazolyl, or pyridyl in which the phenyl may bear one or two substituents independently selected from halo, cyano, hydroxy, methoxy, acetoxy, benzyloxy, amino, acetylamino,  
20 nitro and 3,4-methylenedioxy, and the thienyl or furyl may bear a methyl or nitro substituent);

or a prodrug of the compound of formula I;

or a pharmaceutically acceptable salt of the compound of formula I or prodrug thereof.

25

15. The compound of Claim 14 wherein

A<sup>3</sup>, A<sup>4</sup>, A<sup>5</sup> and A<sup>6</sup>, together with the two carbons to which they are attached, complete a substituted heteroaromatic ring in which

30 (a) one of A<sup>3</sup>, A<sup>4</sup>, A<sup>5</sup> and A<sup>6</sup> is N, and each of the others is CR<sup>3</sup>, CR<sup>4</sup>, CR<sup>5</sup> or CR<sup>6</sup>, respectively;

(b) two adjacent residues of A<sup>3</sup>, A<sup>4</sup>, A<sup>5</sup> and A<sup>6</sup> together form S, and each of the others is CR<sup>3</sup>, CR<sup>4</sup>, CR<sup>5</sup> or CR<sup>6</sup>, respectively;

- 68 -

(c) two non-adjacent residues of A<sup>3</sup>, A<sup>4</sup>, A<sup>5</sup> and A<sup>6</sup> are each N, and each of the others is CR<sup>3</sup>, CR<sup>4</sup>, CR<sup>5</sup> or CR<sup>6</sup>, respectively; or

(d) A<sup>3</sup> and A<sup>4</sup> together form a fused benz ring, and A<sup>5</sup> and A<sup>6</sup> together form -NH-;  
 5 wherein

each of R<sup>3</sup>, R<sup>4</sup>, R<sup>5</sup> and R<sup>6</sup> is hydrogen, or one or two of R<sup>3</sup>, R<sup>4</sup>, R<sup>5</sup> and R<sup>6</sup> is independently chloro, bromo or methyl and the others are hydrogen;


10 L<sup>1</sup> is -NH-CO- or -CO-NH- such that -L<sup>1</sup>-Q<sup>1</sup> is -NH-CO-Q<sup>1</sup> or -CO-NH-Q<sup>1</sup>;

Q<sup>1</sup> is phenyl, 2-thienyl, 4-thiazolyl, 2-pyridyl, 2-naphthyl or 1,2-benzisoxazol-6-yl in which the phenyl may bear one, two or three substituents at the 3-, 4- or  
 15 5-position(s) independently selected from halo, cyano, carbamoyl, aminomethyl, methyl, methoxy, hydroxymethyl, formyl, vinyl, amino, hydroxy and 3,4-methylenedioxy, the 2-thienyl may bear a chloro or methyl substituent at the 5-position, the 4-thiazolyl may bear an amino substituent at  
 20 the 2-position, the 2-pyridyl may bear an amino substituent at the 6-position, and the 1,2-benzisoxazol-6-yl may bear a chloro or methyl substituent at the 3-position;

R<sup>2</sup> is -L<sup>2A</sup>-Q<sup>2A</sup>, -L<sup>2B</sup>-Q<sup>2B</sup>, -L<sup>2C</sup>-Q<sup>2C</sup> or -L<sup>2D</sup>-Q<sup>2D</sup> wherein

L<sup>2A</sup> is a direct bond; and

25 Q<sup>2A</sup> is




in which D is carbonyl or -CHR<sup>k</sup>- in which R<sup>k</sup> is hydrogen, hydroxy, (1-6C)alkoxy or -CH<sub>2</sub>-R<sup>j</sup> in which R<sup>j</sup> is carboxy,  
 30 [(1-4C)alkoxy]carbonyl or carbamoyl which may bear one or two (1-2C)alkyl substituents on the nitrogen; and one of R<sup>m</sup>

- 69 -

and  $R^n$  is hydrogen and the other is amino, bromo, (1-4C)alkyl or (1-4C)alkoxy, or  $R^m$  and  $R^n$  together form a benz ring;

$L^{2B}$  is -NH-CO-, -O-CO-, -CH<sub>2</sub>-O- or -O-CH<sub>2</sub>- such that  
 5 - $L^{2B}$ - $Q^{2B}$  is -NH-CO- $Q^{2B}$ , -O-CO- $Q^{2B}$ , -CH<sub>2</sub>-O- $Q^{2B}$  or -O-CH<sub>2</sub>- $Q^{2B}$ ;  
 and

$Q^{2B}$  is



10 in which  $R^o$  is hydrogen, halo, (1-6C)alkyl, (1-4C)alkoxy, benzyloxy or (1-4C)alkylthio; and  $R^p$  is 1-hydroxyethyl, 1-hydroxy-1-methylethyl, 1-methoxy-1-methylethyl, 4-piperidinyl, 4-pyridinyl, dimethylaminosulfonyl or -J- $R^q$  in which J is a single bond, methylene, carbonyl, oxo,  
 15 -S(O)<sub>q</sub>- (wherein q is 0, 1 or 2), or -NR<sup>r</sup>- (wherein R<sup>r</sup> is hydrogen or methyl); and  $R^q$  is (1-6C)alkyl, phenyl, 3-pyridyl or 4-pyridyl;

$L^{2C}$  is -NR<sup>v</sup>-CO-X-, -NR<sup>v</sup>-CS-Y-, -CH<sub>2</sub>-CO-NR<sup>w</sup>-CH<sub>2</sub>-,  
 -O-CO-, -O-CH<sub>2</sub>-, -S-CH<sub>2</sub>- or -CH<sub>2</sub>-NR<sup>x</sup>-CH<sub>2</sub>- such that - $L^{2C}$ - $Q^{2C}$   
 20 is -NR<sup>v</sup>-CO-X- $Q^{2C}$ , -NR<sup>v</sup>-CS-Y- $Q^{2C}$ , -CH<sub>2</sub>-CO-NR<sup>w</sup>-CH<sub>2</sub>- $Q^{2C}$ ,  
 -O-CO- $Q^{2C}$ , -O-CH<sub>2</sub>- $Q^{2C}$ , -S-CH<sub>2</sub>- $Q^{2C}$  or -CH<sub>2</sub>-NR<sup>x</sup>-CH<sub>2</sub>- $Q^{2C}$  in  
 which X is -(CH<sub>2</sub>)<sub>x</sub>- (wherein x is 0, 1 or 2), -NR<sup>w</sup>-CH<sub>2</sub>-,  
 -O-CH<sub>2</sub>- or -S-CH<sub>2</sub>-; Y is -NR<sup>w</sup>-CH<sub>2</sub>- or -O-CH<sub>2</sub>-; each of R<sup>v</sup>  
 and R<sup>w</sup> is independently hydrogen, benzyl or (1-6C)alkyl  
 25 which is not branched at the α-position; and R<sup>x</sup> is hydrogen, benzyloxycarbonyl or [(1-4C)alkoxy]carbonyl; and

$Q^{2C}$  is 1-(4-pyridyl)piperidin-4-yl in which the pyridyl may bear a substituent at its 2-position selected from cyano, aminomethyl, carboxy, hydroxymethyl and (1-2C)alkyl;

30  $L^{2D}$  is -NH-CO- such that - $L^{2D}$ - $Q^{2D}$  is -NH-CO- $Q^{2D}$ ; and  
 $Q^{2D}$  is selected from 4-(4-pyridinyl)benzyloxy, 9-oxo-9H-fluoren-3-yl, benzo[b]thiophen-2-yl (which may bear a

- 70 -

chloro, methyl or methoxy substituent), benzofuran-2-yl (which may bear a chloro, methyl or methoxy substituent), 4-(4-morpholinyl)-4-oxobutyl, and 4-piperidinyl bearing a substituent at the 1-position selected from methylsulfonyl, phenylsulfonyl and  $-\text{CH}_2-\text{R}^Z$  in which  $\text{R}^Z$  is isopropyl, cyclopropyl, phenyl, furyl, thienyl, 2-thiazolyl, or pyridyl in which the phenyl may bear one or two substituents independently selected from halo, cyano, hydroxy, methoxy, acetoxy, benzyloxy, amino, acetylamino, nitro and 3,4-methylenedioxy, and the thienyl or furyl may bear a methyl or nitro substituent;

or a prodrug of the compound of formula I;

or a pharmaceutically acceptable salt of the compound of formula I or prodrug thereof.

15

16. The compound of Claim 14 or 15 wherein for an alkyl group or the alkyl portion of an alkyl containing group, (1-2C)alkyl is methyl or ethyl; (1-4C)alkyl is methyl, ethyl, propyl, isopropyl, butyl, isobutyl, or t-butyl; (1-6C)alkyl is methyl, ethyl, propyl, butyl, pentyl or hexyl; and halo is bromo or chloro.

17. The compound of Claim 16 wherein for an alkyl group or the alkyl portion of an alkyl containing group, (1-2C)alkyl is methyl; (1-4C)alkyl is methyl, isopropyl, butyl or t-butyl; (1-6C)alkyl is methyl, butyl or hexyl; and halo is chloro.

18. The compound of any of the above Claims 14-17 wherein  $\text{R}^2$  is selected from  $-\text{L}^{2\text{A}}-\text{Q}^{2\text{A}}$ ,  $-\text{NH}-\text{CO}-\text{Q}^{2\text{B}}$ ,  $-\text{NRV}-\text{CO}-\text{X}-\text{Q}^{2\text{C}}$ ,  $-\text{NRV}-\text{CS}-\text{Y}-\text{Q}^{2\text{C}}$ , and  $-\text{NH}-\text{CO}-\text{Q}^{2\text{D}}$ .

19. The compound of any of the above Claims 14-18 wherein the compound of formula I is a pyridine in which one



- 71 -

of A<sup>3</sup>, A<sup>4</sup>, A<sup>5</sup> and A<sup>6</sup> is N, and each of the others is CR<sup>3</sup>, CR<sup>4</sup>, CR<sup>5</sup> or CR<sup>6</sup>, respectively.

20. The compound of any of the above Claims 14-19  
5 wherein the compound of formula I is a thiophene in which the two adjacent residues A<sup>5</sup> and A<sup>6</sup> together form S, and A<sup>3</sup> and A<sup>4</sup> are CR<sup>3</sup> and CR<sup>4</sup>, respectively.

21. The compound of any of the above Claims 14-19  
10 wherein the compound of formula I is an indole in which the two adjacent residues A<sup>5</sup> and A<sup>6</sup> together form -NH-, and A<sup>3</sup> and A<sup>4</sup> together form a fused benz ring.

22. The compound of Claim 19 wherein A<sup>4</sup> is N, and each  
15 of R<sup>3</sup>, R<sup>5</sup> and R<sup>6</sup> is hydrogen.

23. The compound of any of the above Claims 14-22  
wherein Q<sup>1</sup> is 4-methoxyphenyl.

20 24. The compound of any of the above Claims 14-23 wherein R<sup>2</sup> is (4-t-butylbenzoyl)amino, (4-methoxybenzoyl)-amino, [4-(4-pyridyl)benzoyl]amino or [1-(4-pyridyl)-piperidin-4-yl]methoxycarbonylamino.

25 25. The compound of any of the above Claims 14-24 wherein L<sup>1</sup>-Q<sup>1</sup> is -NH-CO-Q<sup>1</sup>.


26. The compound of any of the above Claims 14-24  
30 wherein L<sup>1</sup>-Q<sup>1</sup> is -CO-NH-Q<sup>1</sup>.

27. A pharmaceutical composition comprising a compound of formula I, or prodrug or pharmaceutically acceptable salt thereof, as claimed in Claim 14 in association with a pharmaceutically acceptable carrier, excipient or diluent.

- 72 -

28. A process for preparing a novel compound of formula I (or a pharmaceutically acceptable salt thereof) as provided in Claim 14 which is selected from


- 5 (A) for a compound of formula I in which the linkage of  $R^2$  to the ring terminates in  $-NH-CO-$ ,  $-NR^V-CO-$  or  $-NR^V-CS-$ , acylating an amine of formula II,



10

or a corresponding amine in which the nitrogen bears the group  $R^V$ , using a corresponding acid which terminates with the group  $HO-CO-$  or  $HO-CS-$ , or an activated derivative thereof;

- 15 (B) for a compound of formula I in which  $-L^1-Q^1$  is  $-NH-CO-Q^1$ , acylating an amine of formula III




- 20 using an acid of formula  $HO-CO-Q^1$ , or an activated derivative thereof;

(C) for a compound of formula I in which  $-L^1-Q^1$  is  $-CO-NH-Q^1$  and  $R^2$  is of the form  $-NH-CO-Q^2$ , acylating an amine of formula  $H_2N-Q^1$  using a [1,3]oxazine of formula IV,


25

- 73 -



wherein  $Q^2$  represents  $Q^{2B}$ ,  $Q^{2C}$  or  $Q^{2D}$ ;

- (D) for a compound of formula I in which  $R^2$  is  
 5 - $L^{2A}$ - $Q^{2A}$  and D is carbonyl, diacylating a compound of  
 formula II using an anhydride of formula V;



- whereafter, for any of the above procedures, when a  
 10 functional group is protected using a protecting group,  
 removing the protecting group;

- whereafter, for any of the above procedures, when a  
 pharmaceutically acceptable salt of a compound of formula I  
 is required, it is obtained by reacting the basic form of a  
 15 basic compound of formula I with an acid affording a  
 physiologically acceptable counterion or the acidic form of  
 an acidic compound of formula I with a base affording a  
 physiologically acceptable counterion or by any other  
 conventional procedure; and

- 20 wherein, unless otherwise specified,  $L^1$ ,  $Q^1$ ,  $R^2$ ,  $R^m$ ,  
 $R^n$ ,  $A^3$ ,  $A^4$ ,  $A^5$  and  $A^6$  have any of the values defined in  
 Claim 14.

- 25 29. The use of a factor Xa inhibiting compound of  
 formula I substantially as hereinbefore described with  
 reference to any of the Examples.

- 74 -

30. A novel compound of formula I substantially as hereinbefore described with reference to any of the Examples.

5

31. A process for preparing a novel compound of formula I substantially as hereinbefore described with reference to any of the Examples.

## INTERNATIONAL SEARCH REPORT

International application No.

PCT/US98/13384

<b>A. CLASSIFICATION OF SUBJECT MATTER</b> IPC(6) :Please See Extra Sheet. US CL :Please See Extra Sheet. According to International Patent Classification (IPC) or to both national classification and IPC		
<b>B. FIELDS SEARCHED</b> Minimum documentation searched (classification system followed by classification symbols) U.S. : 514/255, 256, 332, 352, 419, 447; 544/325, 358, 407; 546/265, 308; 548/483; 549/69 Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched Electronic data base consulted during the international search (name of data base and, where practicable, search terms used) CAS ONLINE		
<b>C. DOCUMENTS CONSIDERED TO BE RELEVANT</b>		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	UCHIDA et al. Reactions of n-acylaminoacetamide with 1, 3-bifunctional compounds. Bulletin of the Chemical Society of Japan. October 1973, Vol. 46, No. 10, pages 3177-3280, especially 3278, compound VI.	1-5, 14-17, 27, 28
A	WALLIS R. B. Inhibitors of coagulation factor Xa: from macromolecular beginnings to small molecules. Current Opinion in Therapeutic Patents. August 1993, Vol. 3, No. 8, pages 1173-1179.	1-5, 14-17, 27, 28
<input checked="" type="checkbox"/> Further documents are listed in the continuation of Box C. <input type="checkbox"/> See patent family annex.		
* Special categories of cited documents: *A* document defining the general state of the art which is not considered to be of particular relevance *B* earlier document published on or after the international filing date *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified) *O* document referring to an oral disclosure, use, exhibition or other means *P* document published prior to the international filing date but later than the priority date claimed	*T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art *A* document member of the same patent family	
Date of the actual completion of the international search 04 SEPTEMBER 1998		Date of mailing of the international search report 14 OCT 1998
Name and mailing address of the ISA/US Commissioner of Patents and Trademarks Box PCT Washington, D.C. 20231 Facsimile No. (703) 305-3230		Authorized officer <i>Evelyn Huang</i> EVELYN HUANG Telephone No. (703) 308-1235

## INTERNATIONAL SEARCH REPORT

International application No.  
PCT/US98/13384

## C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	Chem. abstr., Vol. 99, No. 23, 05 December 1983 (Columbus, OH, USA) page 743, column 2, the abstract No. 194839t, LIU et al. 'Potential antineoplastic sulphydryl agents. IV. Synthesis of N1-phenylindole-2,3-dicarboxamide.' Tai-wan Yao Hsueh Tsa Chih. 1983, 3591), 102-4 (Eng).	1-5, 14-17, 27, 28
A	Chem. abstr., Vol. 118, No. 9, 01 March 1993 (Columbus, OH, USA), page 824, column 1, the abstract No. 80809p, ISHIKAWA et al. 'Preparation of 2-acylpyridine derivatives as agrochemical fungicides.' Jpn. Kokai Tokkyo Koho JP 04,187,675, 06 July 1992.	1-5, 14-17, 27, 28

## INTERNATIONAL SEARCH REPORT

International application No.

PCT/US98/13384

**Box I Observations where certain claims were found unsearchable (Continuation of Item 1 of first sheet)**

This international report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:  
because they relate to subject matter not required to be searched by this Authority, namely:
2. ☐ Claims Nos.:  
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:
3. ☒ Claims Nos.: 6-13, 18-26, 29-31  
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

**Box II Observations where unity of invention is lacking (Continuation of Item 2 of first sheet)**

This International Searching Authority found multiple inventions in this international application, as follows:

Please See Extra Sheet.

1. ☒ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

☐

The additional search fees were accompanied by the applicant's protest.

☐

No protest accompanied the payment of additional search fees.

# INTERNATIONAL SEARCH REPORT

International application No.  
PCT/US98/13384

## A. CLASSIFICATION OF SUBJECT MATTER:

IPC (6):

A 61K 31/38, 31/40, 31/44, 31/495, 31/505; C07D 209/30, 209/40, 213/75, 213/81, 213/82, 239/32, 241/18, 241/20, 333/06, 401/10, 401/12

## A. CLASSIFICATION OF SUBJECT MATTER:

US CL :

514/255, 256, 332, 352, 419, 447; 544/325, 358, 407; 546/265, 308; 548/483; 549/69

## BOX II. OBSERVATIONS WHERE UNITY OF INVENTION WAS LACKING

This ISA found multiple inventions as follows:

This application contains the following inventions or groups of inventions which are not so linked as to form a single inventive concept under PCT Rule 13.1. In order for all inventions to be searched, the appropriate additional search fees must be paid.

Group I, claim(s) 6, 9, 19, 22 and claims 1-5, 10-18, 23-31 in part, drawn to a pyridine compound.

Group II, claim(s) 7, 20 and claims 1-5, 10-18, 23-31 in part, drawn to a thiophene compound.

Group III, claim(s) 8, 21 and claims 1-5, 10-18, 23-31 in part, drawn to an indole compound.

Group IV, claims 1-5, 10-18, 23-31 in part, drawn to a compound of formula I wherein two non-adjacent residues of A3, A4, A5 and A6 are each N.

The inventions listed as Groups I-IV do not relate to a single inventive concept under PCT Rule 13.1 because, under PCT Rule 13.2, they lack the same or corresponding special technical features for the following reasons: group I is drawn to a monocyclic six-membered ring containing one nitrogen, group II is drawn to a monocyclic, sulphur containing 5-membered ring, group III is drawn to a bicyclic nitrogen containing compound whereas group IV is drawn to a monocyclic 6-membered ring containing 2 nitrogen atoms. A pyridine, a thiophene, an indole and a pyrimidine or pyrazine would not have been of sufficient similarity to allow for a Markush grouping to exhibit utility, absent some teaching of equivalence in the prior art.